

Study Analysis Group 15
Exoplanet Exploration Program Analysis Group

Science Questions for Direct Imaging Missions

DRAFT



Contact: Daniel Apai (apai@arizona.edu)

March 19, 2016

Table of Contents

Table of Contents	3
SAG15 Membership	5
Introduction	6
Overview of science questions	7
Discussion of Science Questions	8
A1. What is the diversity of planetary architectures? Are there typical classes/types of planetary architectures?	8
A2. What are the distributions and properties of planetesimal belts and eco-zodiacal disks in exoplanetary systems and what can these tell about the formation and dynamical evolution of the planetary systems?	9
A3. Are there natural classes of exoplanets and if so, what are these?	10
A4. How Common is the Solar System?	11
A5. What are typical formation/evolution pathways for planetary systems?	11
Category B: Questions on Exoplanet Properties	12
B1. How do rotational periods and obliquity vary with orbital elements and planet mass/type?	12
B2. Which rocky planets have liquid water on their surfaces?	13
B3. What are the origins and composition of clouds and hazes and how do these vary with system parameters?	14
B4. Which planets have large continents and oceans?	14
Category C Questions: Processes	15
C1. What processes/properties set the modes of atmospheric circulation and heat transport in exoplanets and how do these vary with system parameters?	15
C2. What are the Key Evolutionary Pathways for Rocky Planets?	15
C3. What types/which planets have active geological activity, interior processes, and / or continent-forming/resurfacing processes?	16
Data Requirements	17
Summary of data requirements	17
SAG15 Charter	18
SAG15 Timeline and Process	20

SAG15 Membership

Chair: Daniel Apai, University of Arizona (apai@arizona.edu)

Members:

Travis Barman, University of Arizona
Alan Boss, Carnegie DTM
James Breckenridge, Caltech
David Ciardi, IPAC/Caltech
Ian Crossfield, University of Arizona
Nicolas Cowan, McGill University
William Danchi, NASA GSFC
Shawn Domagal-Goldman, NASA GSFC
Caroline Morley, Lick Observatory
Glenn Schneider, University of Arizona
Nicolas Iro, University of Hamburg
Stephen Kane, San Francisco State University
Theodora Karalidi, University of Arizona
James Kasting, Penn State University
Ravikumar Kopparapu, NASA GSFC

Patrick Lowrence, IPAC/Caltech
Avi Mandell, NASA GSFC
Mark Marley, NASA Ames
Michael McElwain, NASA GSFC
Nikku Madhusudhan, Cambridge University
Charley Noecker, JPL
Peter Plavchan, Missouri State University
Aki Roberge, NASA GSFC
Leslie Rogers, University of Chicago
Adam Showman, University of Arizona
Arif Solmaz
Philip Stahl, NASA MSFC
Karl Stapelfeldt, JPL
Mark Swain, JPL
Margaret Turnbull, SETI Institute

SAG15 Website: <http://eos-nexus.org/sag15>

Introduction

This report presents organized input from the international exoplanet community on science questions that can be answered by direct imaging missions.

For each science question we also explore the types and quality of datasets that are either required to answer the question or greatly enhance the quality of the answer. We also highlight questions that require or benefit from complementary (non-direct imaging) observations.

In preparing the report no specific mission architecture or requirements were assumed or advocated for; however, where obvious connections to planned or possible future mission existed there were identified. The report does not include discussion of biosignatures or planets transformed by life; but it does include discussion of the characterization of habitable zone earth-sized planets.

Overview of science questions

Science Questions on Exoplanetary System Architectures & Population	Importance
A1. What is the diversity of planetary architectures? Are there typical classes/ types of planetary architectures?	
What are the distributions and properties of planetesimal belts and eco-zodiacal disks in exoplanetary systems and what can these tell about the formation and dynamical evolution of the planetary systems?	
A3. Are there natural classes of exoplanets and if so, what are these?	
A4. How Common is the Solar System?	
A5. What are typical formation/evolution pathways for planetary systems?	
Science Questions on Exoplanet Properties	Importance
B1. How do rotation periods and obliquity vary with orbital elements and planet mass/type?	
B2. Which rocky planets have liquid water on their surfaces?	
B3. What are the origins and composition of clouds and hazes and how do these vary with system parameters?	
B4. Which planets have large continents and oceans?	
Science Questions of Evolution and Processes that Change Exoplanets	Importance
C1. What processes/properties set the modes of atmospheric circulation and heat transport in exoplanets and how do these vary with system parameters?	
C2. What are the Key Evolutionary Pathways for Rocky Planets?	
C3. What types/which planets have active geological activity, interior processes, and /or continent-forming/resurfacing processes?	

Discussion of Science Questions

The following sections provide a preliminary discussion of possible science questions for direct imaging missions. The questions are organized in three categories: Questions in Category A focus on the statistical assessment of the properties of exoplanetary systems. Questions in Category B focus on the understanding of specific exoplanets. Questions in Category C focus on understanding the details and importance of key processes by establishing causal relations between present-day properties and processes.

A1. What is the diversity of planetary architectures? Are there typical classes/types of planetary architectures?

Our current picture of planetary system architectures builds on five sources: 1) Solar System; 2) Data from transiting exoplanets, primarily Kepler, which probes the inner planetary systems (typically up to periods of approximately 1 year); 3) radial velocity surveys, which provide data on planets with masses typically larger than those accessible to Kepler observations, but some of which cover multi-year periods; 4) microlensing surveys, which are also sensitive to small rocky planets at intermediate periods, but provide a yet limited statistics; 5) direct imaging surveys: capable of probing giant exoplanets at semi-major axes of 8 au or longer.

Based on the extrapolation of the close-in exoplanet population detected by Kepler we do not yet have an efficient method to detect most of the planets (at intermediate to large periods, with masses comparable to Earth).

A direct imaging mission could provide a powerful tool to survey planets at intermediate orbits (~1 to 30 au), establish their orbits or constrain their orbital parameters, and measure or deduce their masses or sizes.

Comments from Telecon 2:

SDG: There will be an upcoming white paper by Debra Fischer and colleagues on high-precision RV measurements that will be very relevant for this question.

SDG: Rather than identifying a complete understanding of planetary architectures (which may be impossible to achieve), we may want to identify specific hypotheses that should be tested. For example, a hypothesis on the transition between rocky planets and gas giants could have specific, testable predictions.

Data Required: *Optical or infrared imaging to identify the presence and location of planets in each system. Multi-epoch imaging (or complementary radial velocity or astrometry) is required to constrain orbital parameters.*

Questions to SAG15: 1) *How many epochs are required to establish orbital parameters?*
2) *To what accuracy should the orbital parameters be measured to?*

3) *What sample size (number of systems imaged) would be a) minimum required, or be b) optimally suited for answering this question?*

Complementary Non-Imaging Data:

- 1) Radial velocity: Constraints from radial velocity measurements can greatly reduce the number of direct imaging epochs required to establish the orbital elements of the planets. These measurements can also constrain or determine the mass of the target planets.
- 2) Astrometry: Constraints from radial velocity measurements can greatly reduce the number of direct imaging epochs required to establish the orbital elements of the planets. These measurements can also constrain or determine the mass of the target planets.
- 3) Ground-based adaptive optics imaging: These observations may be capable of discovering giant exoplanets and providing positions at additional epochs.

A2. What are the distributions and properties of planetesimal belts and exozodiacal disks in exoplanetary systems and what can these tell about the formation and dynamical evolution of the planetary systems?

Direct imaging missions will likely provide spatially resolved images of exo-zodi disks, possibly composed of narrow and/or extended dust belts. In these belts dust is produced by minor body collisions and the dust belts are dynamically sculpted by the gravitational influence of the planets in the systems, grain-grain collisions, as well as radiation pressure from the stars.

Currently, detailed simulations of debris disk structures and disk-planet interactions provide predictions for the expected disk structures. In a large set of disks complex structures have been observed which have been explained via the influence of yet unseen planets; in a very small set of systems disks and planets have been observed together, providing an opportunity to study disk-planet interactions and to validate models.

Questions to SAG15:

- 1) What are the most compelling science questions to be answered through observations of dust belts / exozodiacal disk structures?
- 2) To what extent could the dust belt structures be used to: a) deduce the presence of lower-mass planets; b) provide constraints on the mass and eccentricity of the directly imaged planets in the system; or c) constrain the dynamical evolution of given planetary systems, i.e., through constraining possible migration histories?

A3. Are there natural classes of exoplanets and if so, what are these?

For centuries the planets in the Solar System provided a basic template for planet classification, based on their mass distribution and composition (“rocky planets”, “gas giants”, “ice giants”, “dwarf planets”). In the Solar System the mass/composition based classification also correlates with an alternative location-based definition and agrees with differences in the expected formation pathways (i.e., accretion of rock/iron; gas accretion to a rocky/icy core; gas-limited accretion to a rocky/icy core).

However, the diversity of exoplanets discovered revealed a greater diversity of exoplanets, including an apparently continuous planetary mass distribution, presence of planets of different sizes/compositions at very different orbits, and possibly multiple very different formation and evolutionary pathways.

Publications converged to ad hoc planet categories and adjectives (e.g., *by mass*: sub-earths, earths, super-earths, mini-neptunes, neptunes, saturns, jupiters, super-jupiter; *by temperature*: cold, warm, hot; *by formation*: exoplanet, planetary-mass companion, brown dwarf, ejected planet).

Planet formation and evolution pathways may lead to continuous spectra in planet mass, composition, temperature, and formation/evolution, in which case it is of limited use (or may be misleading) to distinguish different categories of exoplanets. On the other hand it is also possible or even likely that in a multi-dimensional space of planet properties strong clustering is present, providing natural new categories for planets.

Identifying the natural categories of planets can enhance the effectiveness of studies (by comparing similar objects).

Imaging Data Requirements:

Complementary Non-Imaging Data:

- 1) Radial velocity / Astrometry: Required for mass determination.
- 2) Transits: Statistics on short-period planets not accessible to direct imaging.

Comments on Telecon 2:

SDG: We should also consider that, given also complementary measurements, which dimensions of the description of planetary architectures are important and which are not probed sufficiently by existing/future measurements, i.e., prioritize dimensions that are important and unexplored.

SK: We should consider dynamical constraints, such as mean-motion resonances, on the N-dim parameter space, as the dynamical information can be important for understanding the systems.

SK: A related question: are compact planetary systems extremely common? There are some biases and intriguing trends in the existing transit-based data that could be verified with direct imaging.

A4. How Common is the Solar System?

Needs more discussion:

The Solar System may in many ways differ from typical planetary systems. Which parameters are relevant for this question?

Comments from Telecon 2:

SK? : Jupiter analogs are becoming detectable and this question (at least frequency of Sun-Jupiter pairs) will be answered relatively soon.

DA: There are many parameters that could be used to determine the similarity of a given planetary system and the Solar System; not clear which are more important than others for our purposes and how to weight them.

Consensus: We will need to better define this question.

Complementary Observations:

Type of Data:

A5. What are typical formation/evolution pathways for planetary systems?

Suggested toward at telecon 2 - needs more discussion.

Complementary Observations.

Type of Data:

Category B: Questions on Exoplanet Properties

B1. How do rotational periods and obliquity vary with orbital elements and planet mass/type?

Rotational Periods: Rotational periods are important parameters of planets as they constrain: 1) atmospheric dynamics; 2) diurnal surface temperature variations; and 3) formation and angular evolution.

For potentially habitable planets rotation rates are an important parameter in climate models. Current calculations assume a rotational rate that matches Earth's rate.

Simulated and actual observations of rocky and giant exoplanets demonstrate that time-resolved precision photometry or spectroscopy can be used to directly measure the rotational periods of exoplanets (references). Ongoing observations by the Earth-observing EPIC camera (Kane et al.) will provide an important demonstration of the technique.

Obliquity: The obliquity of habitable planets has a major impact on the seasonal and diurnal temperature variations and on their climate in general. Obliquity is much more difficult to determine than the directly observable rotational rate. However, simulated observations demonstrate that it is possible to determine this quantity from high signal-to-noise reflected light lightcurves obtained at multiple orbital phases.

Questions to SAG15:

- 1) *How important are rotation periods for different types of planets?*
- 2) *To what accuracy should rotation periods be determined?*

Comments from Telecon 2:

RK: Not clear how these could be measured.

DA: Rotation periods could be directly measured from time-resolved precision photometry. Obliquity is much more difficult, but was shown that it could be constrained from the combination of orbital phase and rotational phase maps.

SK: We are carrying out a study using the Earth-observing EPIC camera to study Earth as exoplanet. We will also demonstrate how rotational rate can be determined and what other parameters can be constrained.

RK: Rotation rate is also important for habitability calculations; most current calculations just assume Earth-like rotation rate.

Consensus: Useful question.

B2. Which rocky planets have liquid water on their surfaces?

Water is not a biosignature itself, but the presence of liquid water is required for life as we know it. And life as we know it is probably the only kind of life that we may be able to identify remotely. Liquid water is not the only factor required for planetary habitability, but it is arguably the most important one.

The presence of liquid surface water may be constrained through different measurements, including atmospheric spectroscopy; rotational phase mapping; presence of clouds composed of liquid water droplets.

Liquid water is a “habitability signature”. Establishing which habitable zone planets have liquid water on their surfaces provides an important context for EXOPAG SAG16, which focuses on biosignatures, but will rely on SAG15 for habitability signatures and characterization of habitable planets.

Comments from SAG15 Team / Telecon 2:

From Jim Kasting:

The key question for me concerns the availability of liquid water on the surfaces of rocky planets orbiting other stars. Water is not a biosignature itself, but the presence of liquid water is required for life as we know it. And life as we know it is probably the only kind of life that we may be able to identify remotely. Liquid water is not the only factor required for planetary habitability, but it is arguably the most important one.

Related observables: 1) orbital distance. It is critical to determine the semi-major axes of rocky exoplanets because this plays a key role in determining whether they can support liquid water on their surfaces. Note that this likely requires multiple revisits to each planetary system of interest. On TPF-C, we assumed 5-6 revisits per system. 2) Gaseous CO₂ and H₂O. Both of these gases are potentially observable at near-IR wavelengths. CO₂ is even easier to observe in the thermal-IR, but that requires a different kind of direct imaging mission.

Other comments:

Consensus: Important question.

Multiple people: Presence of water may be constrained through measurements other than atmospheric spectroscopy; rotational phase mapping, for example, could be important.

RK: Determining P-T profile and presence of water clouds would also be important/ useful.

SDG: This question connects well to SAG16, where we study biosignatures. Liquid water on the planetary surface is a habitability signature, which is not studied in SAG16; SAG16 will rely on SAG15 for this context.

B3. What are the origins and composition of clouds and hazes and how do these vary with system parameters?

All Solar System planets with an atmosphere harbor cloud and/or haze layers. Clouds and hazes influence the pressure-temperature structure of the atmosphere, its emission and transmission spectrum, the albedo of the planets. Clouds and hazes may also be used as tracers to characterize atmospheric dynamics (circulation, mixing, turbulence).

Presence of haze or cloud layers may also mask the presence of specific atmospheric absorbers even if present at large abundances at pressures higher than the particle layer.

Exoplanets are expected to harbor a large variety of condensates and haze-forming particles that may be important and/or observable with future direct imaging missions.

Understanding the composition of cloud- and haze-forming particles is an important step in developing physical/chemical models for exoplanets.

Questions to SAG15:

- 1) What data can constrain particle size distribution, pressure levels, composition?
- 2) What fundamental parameters (composition, temperature, surface gravity?) are expected to have significant impact on cloud/haze formation and properties?

B4. Which planets have large continents and oceans?

Analysis of simulated eco-earth observations was used to demonstrate that rotational phase mapping (time-resolved observations of hemisphere-integrated reflected light from the planet) can reveal the types and distribution of surfaces. Equipped with additional data on the color/spectra of the features and the physical conditions on the planetary surface may be used to identify surface features as oceans and continents.

Questions to SAG15:

Category C Questions: Processes

C1. What processes/properties set the modes of atmospheric circulation and heat transport in exoplanets and how do these vary with system parameters?

Atmospheric circulation plays a key role in redistribution energy in exoplanets' atmospheres. Depending on wind speed, rotational velocity, insolation, latent heat released during condensation, and other system parameters a range of atmospheric circulation regimes are expected on planets that can be studied with direct imaging missions.

For potentially habitable exoplanets atmospheric circulation will determine the day-night heat differential and the equator-pole temperature difference. Understanding the presence and size of Hadley cells can also provide important insights into how water vapor (or other condensibles) may be distributed in habitable planets.

Understanding atmospheric circulation in habitable exoplanets is an important component in establishing a correct climate model for them.

As of now, atmospheric circulation has been probed in the Solar System planets and a small sample of hot jupiters and brown dwarfs. Lower-mass exoplanets remain unexplored.

Using time-resolved observations and rotational phase mapping techniques atmospheric circulation may be constrained to some level.

Questions to SAG15:

- 1) To what level can the atmospheric circulation be constrained for different types of planets?
- 2) What hypotheses / toy circulation models should be tested for gas giants?
- 3) What hypotheses / toy circulation models should be tested for habitable super-earths / earths?
- 4) What data type and cadence is required or best suited for characterizing circulation?

C2. What are the Key Evolutionary Pathways for Rocky Planets?

From Nick Cowan: [WHY DO PLANETS TURN OUT THE WAY THEY DO?](#) In other words, which aspects of planet formation (mass, spin, composition) impact which aspects of planetary atmospheres and climate? This is a classic nature vs nurture problem. Maybe the most important determinant of a planet's evolution is the final major impact, or its star's UV flux, or maybe it is the planet's volatile content... or maybe it is first-order stochastic with only second-order correlations to the factors above. Another way to think about the effect of nurture is the extent to which planets exhibit hysteresis. If Earth and Venus traded locations, would the new cooler Venus end up looking more Earth-like in the long run, and vice versa.

C3. What types/which planets have active geological activity, interior processes, and /or continent-forming/resurfacing processes?

Planetary interior processes and geological activity play an important role in coupling Earth's atmosphere to its crust and providing a long-term stabilizer for Earth's climate. Developing reliable climate models to determine the habitability of potentially habitable planets will likely require assumptions on geological activity and the level of coupling between the planet's crust and atmosphere.

Interior processes are obviously very difficult to probe via low signal-to-noise and spatially unresolved remote sensing.

Possible ideas includes:

- 1) Detection of atmospheric absorbers that can be attributed to volcanic release, and deducing the level of volcanic activity.
- 2) Using land-mass distribution to place constraints on continent-forming processes.
- 3) Other methods?

Data Requirements

This section identifies the type and quality of data ideal or required to answer the individual science questions discussed in Section 2.

Summary of data requirements

Science Questions	Optical			Near-Infrared			Mid-IR		Num. Targets
	Phot.	Spec.	Pol.	Phot.	Spec.	Pol.	Phot.	Spec.	
A1									
A2									
A3									
A5									
B1									
B2									
B3									
B4									
C1									
C2									
C3									
C4									

SAG15 Charter

Future direct imaging missions may allow observations of flux density as a function of wavelength, polarization, time (orbital and rotational phases) for a broad variety of exoplanets ranging from rocky sub-earths through super-earths and neptunes to giant planets. With the daunting challenges to directly imaging exoplanets, most of the community's attention is currently focused on *how* to reach the goal of exploring habitable planets or, more specifically, how to search for biosignatures.

Arguably, however, most of the exoplanet science from direct imaging missions will not come from biosignature searches in habitable earth-like planets, but from the studies of a much larger number of planets *outside* the habitable zone or from planets within the habitable zone that do not display biosignatures. These two groups of planets will provide an essential context for interpreting detections of possible biosignatures in habitable zone earth-sized planets.

However, while many of the broader science goals of exoplanet characterization are recognized, there has been no systematic assessment of the following two questions:

1) What are the most important science questions in exoplanet characterization *apart* from biosignature searches?

2) What type of data (spectra, polarization, photometry) with what quality (resolution, signal-to-noise, cadence) is required to answer these science questions?

We propose to form SAG15 to identify the key questions in exoplanet characterization and determine what observational data obtainable from direct imaging missions is necessary and sufficient to answer these.

The report developed by this SAG will explore high-level science questions on exoplanets ranging from gas giant planets through ice giants to rocky and sub-earth planets, and — in temperatures — from cold (~200 K) to hot (~2,000 K). For each question we will study and describe the type and quality of the data required to answer it.

For example, the SAG15 could evaluate what observational data (minimum sample size, spectral resolution, wavelength coverage, and signal-to-noise) is required to test that different formation pathways in giant planets lead to different abundances (e.g. C/O ratios). Or the SAG15 could evaluate what photometric accuracy, bands, and cadence is required to identify continents and oceans in a habitable zone Earth-sized or a super-earths planet. As another example, the SAG15 could evaluate what reflected light data is *required* to constrain the fundamental parameters of planets, e.g. size (distinguishing earth-sized planets from super-earths), temperature (cold/warm/hot), composition (rocky, icy, gaseous), etc.

SAG15 will not attempt to evaluate exoplanet detectability or specific instrument or mission capabilities; instead, it will focus on evaluating the *diagnostic power* of different measurements on key exoplanet science questions, simply adopting resolution, signal-to-noise, cadence, wavelength coverage as parameters along which the diagnostic power of the data will be studied. Decoupling instrumental capabilities from science goals allows this community-based effort to explore the science goals for exoplanet characterization in an unbiased manner and in a depth beyond what is possible in a typical STDT.

EXOPAG SAG15 Draft Report – Feb 5, 2016

We envision the SAG report to be important for multiple exoplanet sub-communities and specifically foresee the following uses:

- 1) Future STD teams will be able to easily connect observational requirements to missions to fundamental science goals;
- 2) By providing an overview of the key science questions on exoplanets and how they could be answered, it may motivate new, dedicated mission proposals;
- 3) By providing a single, unified source of requirements on exoplanet data in advance of the Decadal Survey, the science yield of various missions designs can be evaluated realistically, with the same set of assumptions.

Our goal is to carry out this SAG study by building on both the EXOPAG and NExSS communities.

We aim to complete a report by Spring 2017 and submit it to a refereed journal, although this timeline can be adjusted to maximize the impact of the SAG15 study for the ongoing and near-future STDTs and other mission planning processes.

Synergy with a potential future SAG proposed by Shawn Domagal-Goldman: While the SAG proposed here will include studies of habitable zone rocky planets, it will focus on planets without significant biological processes. A future SAG may be proposed by Shawn Domagal-Goldman to explore biosignatures; if such a SAG is proposed, we envision a close collaboration on these complementary, but distinct problems.

SAG15 Timeline and Process

This section records the SAG15 timeline and explains how the report was assembled and identifies the methods through which community input was solicited and collected.

SAG15 Website: <http://eos-nexus.org/sag15>