

SAG15 DRAFT REPORT

SCIENCE QUESTIONS FOR DIRECT IMAGING EXOPLANET MISSIONS



DANIEL APAI AND THE SAG15 TEAM

August 16, 2016

Contents

1	The SAG15 Team	3
2	Introduction	4
3	Overview of Science Questions	5
4	Exoplanetary System Characterization	6
4.1	A1. What is the diversity of planetary architectures? Are there typical classes/types of planetary architectures? How common are Solar System-like planetary architectures?	6
4.1.1	Complementary Non-Imaging Data	7
4.2	A2. What are the distributions and properties of planetesimal belts and exozodiacal disks in exoplanetary systems and what can these tell about the formation and dynamical evolution of the planetary systems?	9
4.2.1	Current Knowledge	9
4.2.2	Sub-questions	11
4.2.3	Complementary Data	12
5	Exoplanet Characterization	14
5.1	B1. How do rotational periods and obliquity vary with orbital elements and planet mass/type?	14
5.1.1	State of the Art to Measure Rotational Periods	14
5.1.2	Science Cases	15
5.1.3	The Science Value of Independently Measured Planet Masses and Radii	19
5.2	B2: Which rocky planets have liquid water on their surfaces? Which planets have continents and oceans?	23
5.2.1	Detecting Oceans	23
5.2.2	Inferring Liquid Water	24

5.3	B3. What are the origins and composition of clouds and hazes in ice/gas giants and how do these vary with system parameters?	26
5.3.1	Science Value of Independently Measured Planet Masses and Radii	31
6	Exoplanetary Processes	32
6.1	C1. What processes/properties set the modes of atmospheric circulation and heat transport in exoplanets and how do these vary with system parameters?	32
6.2	C2. What are the key evolutionary pathways for rocky planets and what first-order processes dominate these?	35
6.2.1	Science Value of Independently Measured Planet Masses and Radii	36
6.3	C3. What types/which planets have active geological activity, interior processes, and/or continent-forming/resurfacing processes?	36
6.3.1	Geological Activity and Plate Tectonics on Extrasolar Rocky Planets	38
6.3.2	Observational Methods	39
6.3.3	Complementary Datasets	41
6.3.4	Science Value of Independently Measured Planet Masses and Radii: Very High	42
7	Data Requirements	42
A	SAG15 Charter	43

1. The SAG15 Team

Chair: Daniel Apai, University of Arizona (apai@arizona.edu)

Members:

Travis Barman, University of Arizona	Patrick Lowrence, IPAC/Caltech
Alan Boss, Carnegie DTM	Avi Mandell, NASA GSFC
James Breckenridge, Caltech	Mark Marley, NASA Ames
David Ciardi, IPAC/Caltech	Michael McElwain, NASA GSFC
Ian Crossfield, University of Arizona	Nikku Madhusudhan, Cambridge University
Nicolas Cowan, McGill University	Charley Noecker, JPL
William Danchi, NASA GSFC	Peter Plavchan, Missouri State University
Shawn Domagal-Goldman, NASA GSFC	Aki Roberge, NASA GSFC
Caroline Morley, Lick Observatory	Leslie Rogers, University of Chicago
Glenn Schneider, University of Arizona	Adam Showman, University of Arizona
Nicolas Iro, University of Hamburg	Arif Solmaz
Stephen Kane, San Francisco State University	Philip Stahl, NASA MSFC
Theodora Karalidi, University of Arizona	Karl Stapelfeldt, JPL
James Kasting, Penn State University	Mark Swain, JPL
Ravikumar Kopparapu, NASA GSFC	Margaret Turnbull, SETI Institute

With Additional Input From:

Yuka Fujii, NASA GISS
Markus Kasper, European Southern Observatory
NExSS Community, Breakout session

2. Introduction

This report presents organized input from the international exoplanet community on science questions that can be answered by direct imaging missions.

For each science question we also explore the types and quality of datasets that are either required to answer the question or greatly enhance the quality of the answer. We also highlight questions that require or benefit from complementary (non-direct imaging) observations.

In preparing the report no specific mission architecture or requirements were assumed or advocated for; however, where obvious connections to planned or possible future mission existed these were identified. More detailed evaluations of the capabilities of specific mission architectures are provided in other SAG reports and by ongoing NASA STDTs studies. The SAG15 report does not include discussion of biosignatures or planets transformed by life, which are discussed in the ongoing SAG16 study, however, the SAG15 reports does include discussion of the characterization of habitable zone earth-sized planets.

Community input: Input for this report has been collected and comments on the different report drafts have been solicited through a range of channels, including: i) SAG15 website (<http://eos-nexus.org/sag15>); ii) monthly SAG15 telecons; iii) breakout and discussion sessions during related workshops and meetings; iv) direct requests from topical experts; v) email invitations and solicitations via the EXOPAG and NExSS mailing lists.

Author list and contributor list: The final report will represent the full endorsement of each authors, based on their explicit written statements. In contrast, the SAG15 team list and list of contributors provided in the interim drafts only represents experts who provided input or joined the SAG15 team. The contributor list in the report drafts, therefore, does not represent the endorsement of the draft report and its findings by the contributors.

3. Overview of Science Questions

The science questions in this report are divided into three categories. Questions in Category A aim at the statistical characterization of the formation, evolution, and properties of planetary systems. Questions in Category B aim at the quantitative characterization of individual planets or small groups of planets. Questions in Category C aim at understanding processes that shape planets and planetary atmospheres through comparative studies of planets.

Science Questions on Exoplanetary System Architectures & Population	Importance
A1. What is the diversity of planetary architectures? Are there typical classes/types of planetary architectures? How Common are Planetary Architectures resembling the Solar System?	
A2. What are the distributions and properties of planetesimal belts and eco-zodiacal disks in exoplanetary systems and what can these tell about the formation and dynamical evolution of the planetary systems?	
Science Questions on Exoplanet Properties	Importance
B1. How do rotation periods and obliquity vary with orbital elements and planet mass/type?	
B2. Which rocky planets have liquid water on their surfaces?	
B3. What are the origins and composition of clouds and hazes in ice/gas giants and how do these vary with system parameters?	
B4. How do photochemistry, transport chemistry, surface chemistry, and mantle outgassing effect the composition and chemical processes in terrestrial planet atmospheres (both habitable and non-habitable)?	
Science Questions of Evolution and Processes that Change Exoplanets	Importance
C1. What processes/properties set the modes of atmospheric circulation and heat transport in exoplanets and how do these vary with system parameters?	
C2. What are the Key Evolutionary Pathways for Rocky Planets?	
C3. What types/which planets have active geological activity, interior processes, and /or continent-forming/resurfacing processes?	

4. Exoplanetary System Characterization

4.1. A1. What is the diversity of planetary architectures? Are there typical classes/types of planetary architectures? How common are Solar System-like planetary architectures?

Contributors: Daniel Apai

The term *planetary system architecture* is used here as a descriptor of the high-level structure of a planetary system as given by the stellar mass, the orbits and mass/nature of the planets, and the location/mass of its planetesimal belts.

Understanding the diversity of planetary architectures is important for at least the following two reasons: i) The diversity of planetary system architectures is expected to reflect *the range of possible formation and evolution pathways* of planetary systems. ii) To understand how common are true Earth analogs we must understand *how common are planetary systems with architectures similar to that of the Solar System*.

Our current picture of planetary system architectures builds on five sources: 1) Solar System; 2) Data from transiting exoplanets, primarily the Kepler Space Telescope, which probes the inner planetary systems (typically up to periods of 1 year); 3) radial velocity surveys, which provide data on planets with masses typically larger than those accessible to Kepler observations, but over multi-year periods; 4) microlensing surveys, which are also sensitive to small rocky planets at intermediate periods, but provide a yet limited statistics; 5) direct imaging surveys: capable of probing giant exoplanets at semi-major axes of 8 au or greater.

Based on the extrapolation of the close-in exoplanet population detected by the Kepler mission it is clear that we do not yet have an efficient method to detect the *majority* of exoplanets that exists (at intermediate to large periods, with masses comparable to Earth). ESA's Gaia mission will increase the census of known intermediate- to long-period giant planets by about $\sim 3,000$ new discoveries. In addition, the proper motion information for the Solar neighborhood will improve the identification and age-dating of co-moving stellar groups which, in turn, will greatly reduce the uncertainties in the giant planet mass-to-luminosity conversion used by ground-based direct exoplanet imaging surveys, thereby improving the long-period giant planet occurrence rate and mass distribution measurements.

Furthermore, the gradually extending baselines and improving accuracy of radial velocity measurements will also further improve the occurrence rates for short and intermediate-orbit planets (most significantly for neptune-mass and larger planets). In spite of these

significant improvements the occurrence rates of the sub-neptune planets (including rocky and icy planets) at intermediate- to long-period orbits is presently poorly known and will remain largely unconstrained in the near future.

A direct imaging mission would be powerful in surveying low-mass planets at intermediate and long orbits (1 to 30 au), establishing their orbits or constraining their orbital parameters, and measuring or deducing their masses and sizes.

Although different techniques will sample different planet populations around different set of stars, a capable direct imaging mission will have the unique capability of providing a near-complete census of planets in the targeted systems.

Sub-questions:

- *What is the diversity of planetary architectures?* The statistical assessment of the occurrence rate and mass distribution of planets as a function of system parameters (e.g., stellar mass, composition) can constrain and/or verify planet formation models. The dispersion in different parameters (from data corrected for selection effects and biases) can be used to quantify the diversity of the architectures.
- *Are there typical classes/types of planetary architectures?* If there are different typical planet formation or evolution pathways, these may lead to the emergence of different classes of planetary architectures (e.g., planetary systems with hot jupiters). The presence of classes of planetary systems may be identified as clustering in the multi-dimensional parameter space that describes planetary architectures.
- *How common are Solar System-like planetary architectures?* The *local density* of the systems in the multi-dimensional parameter space that describes planetary architectures, determined at the location of the Solar System provides a measure of the occurrence rate of Solar System-like architectures. Furthermore, in this multi-dimensional parameter space distance-type metrics can be defined to reflect the similarity of any two planetary systems. Although non-unique, such metrics may be used to explore the frequency of systems as a function of distance from the Solar System to establish which nearby systems are the most similar to ours.

4.1.1. Complementary Non-Imaging Data

- *Radial velocity:* Constraints from radial velocity measurements can greatly reduce the number of direct imaging epochs required to establish the orbital elements of the

planets. These measurements can also constrain or determine the mass of the target planets.

- *Microlensing*: Statistical constraints from the WFIRST-Microlensing survey will provide important context for the frequency of medium-separation low-mass planets.
- *Ground-based adaptive optics imaging*: These observations may will be capable of discovering super-earth, neptune, and gas giant exoplanets at intermediate and long periods in nearby systems. By providing positions at additional epochs they will place constraints on the orbits of the planets. In combination with GAIA astrometry the planetary architectures of nearby systems will be relatively well explored in 10-20 years.
- *Gaia Astrometry*: This dataset will provide orbital elements and masses for a large number of intermediate- to long-period gas giant planets, an important statistical context for the planets to be discovered by the direct imaging mission. Furthermore, for individual targets where the direct imaging mission and Gaia can both detect planets, an improved age estimate can be made for the system using giant planet evolutionary models.

Observational Requirements:

Sample size: Medium to large samples (30 to 100).

Observations/Data: Optical or infrared imaging to identify the presence and location of planets in each system. Multi-epoch imaging (or complementary radial velocity or astrometry) is required to constrain orbital parameters.

Pending Questions / Comments:

- 1) How many epochs are required to establish orbital parameters?
- 2) To what accuracy should the orbital parameters be measured to?
- 3) What statistical constraints will WFIRST-ML, Gaia, and future RV surveys provide?
- 4) Add more info on GMT/ELT/TMT capabilities.

4.2. A2. What are the distributions and properties of planetesimal belts and eco-zodiacal disks in exoplanetary systems and what can these tell about the formation and dynamical evolution of the planetary systems?

Contributors: Daniel Apai

Direct imaging missions will provide spatially resolved images of exo-zodiacal disks, possibly composed of narrow and/or extended dust belts. In these belts dust is produced by minor body collisions and the dust belts are dynamically sculpted by the gravitational influence of the star and the planets, grain-grain collisions, as well as radiation pressure (for reviews see, e.g., Wyatt 2008). In some, apparently very rare, systems gas is also present and may introduce the dust distribution.

The distribution and properties of exo-zodiacal dust belts (or debris disks) are important as they provide information on:

- The presence, orbits, and masses of embedded, yet unseen planets.
- The orbits and masses of planets seen in the direct images, but for which orbits are not known.
- The inclination of the disk/planet system.
- The formation and evolution of the system, including the past migration and orbital rearrangements of the planets.
- Compositional constraints on the availability of volatiles/organics in the planetesimal belts and, by inference, in the planets.

4.2.1. Current Knowledge

Currently, large databases of bright debris disks are available for which spatially unresolved thermal infrared observations (spectral energy distributions or SEDs) are available. In addition, for a subset of disks spatially resolved scattered light or thermal emission images




Figure 14. Key structures in the β Pic system, as derived from multi-wavelength imaging.




Fig. 1.— Simulations of the structure of the edge-on debris disk around Beta Pictoris correctly predicted the location and masinteractions of the perturber super-Jupiter Beta Pictoris b (Mouillet et al. 1997). This disk remains the best example of disk-planet. Lower panel: HST/STIS coronagraphic image (blue), ALMA dust continuum (green), and ALMA CO gas emission (red) illustrate the complex structure of the disk (from Apai et al. 2015).

are available (see, e.g., Fig. 1). Mid-infrared spectroscopy of solid state dust features (e.g., Telesco et al. 2005) and polarimetric imaging provide additional constraints on dust composition and disk structure (e.g., Perrin et al. 2015). The different wavelengths, the types of emission (continuum, spectral features), and the polarization properties of the light allow us to disentangle the different dust components and study their origins.

Add a few sentences on general debris disk evolution, collisional events, self-stirring and planetary stirring, orbital rearrangements and bombardments.




Fig. 2.— Comparison of the optical depths predicted by disk-planet interactions models for a composite cloud formed for a 2 earth-mass planet at 6 au (from Stark & Kuchner 2008). The planet, marked with a white dot, orbits counterclockwise in these images. Left: Optical depth of the smallest particles included in the composite clouds. Bottom right: Optical depth of the largest particles included in the composite clouds. The largest particles dominate the optical depth in a cloud of particles released with a Dohnanyi crushing law.

4.2.2. Sub-questions

The presence, orbits, and masses of unseen planets: Detailed simulations of debris disk structures and disk-planet interactions provide predictions for the expected disk structures (see Fig. 2, e.g., Wyatt et al. 1999; Wyatt 2003; Mouillet et al. 1997; Stark & Kuchner 2008). In a large set of disks complex structures have been observed which can possibly be explained by the influence of yet unseen planets (e.g., Schneider et al. 2014); in a very small number of systems disks and planets have been observed together, providing an opportunity to study disk-planet interactions and to validate models (see, e.g., Apai et al. 2015 and Fig. 1).

The orbits and masses of planets seen in the direct images: With certain direct imaging architectures (e.g., starshades) opportunities for multi-epoch observations may be limited, making it more difficult to verify that point sources are planets and not background sources; and to estimate masses/orbits for the planets from short integrations. Most directly imaged systems are expected to host dust disks, whose structures may be used to verify that the planet candidates imaged are indeed in the system and then to constrain their mass and orbit.

The inclination of the disk/planet system: For any planet an important but particularly challenging parameter pair to determine is the inclination/eccentricity pair. These quantities are partially degenerate and can be difficult to disentangle from observations limited to a

handful of visits. Resolved debris disks structures can complement measurements of the planet’s relative motion to break the degeneracy of inclination/eccentricity. For example, nearly-edge on disks can be recognized even in single-epoch images, which then greatly constrain the available parameter space for the planet’s orbit.

The formation and dynamical evolution of the systems: The mass and position of planetesimal belts can provide powerful constraints on the formation and evolution of planetary systems, including planet migration and/or major orbital rearrangements. For example, the asteroid belt and the Kuiper belt in the Solar System have revealed such orbital rearrangement and potential past instabilities (e.g., Malhotra 1993; Tsiganis et al. 2005). In addition, sensitive time-resolved observations in debris disks also have the potential to identify multiple other mechanisms that act on short timescales, such as the aftermath of recent major impacts (e.g., Meng et al. 2014), dust clumps moving under the influence of radiation pressure, or dust created by planetesimals trapped in resonant structures (e.g., Wyatt 2003; Apai et al. 2015; Boccaletti et al. 2015).

Compositional constraints on the availability of volatiles/organics in the planetesimal belts: In each system planetesimal belts are leftover reservoirs of the same material that formed the planets and therefore the planetesimal’s composition constrains the composition of the planets themselves. Of particular interest is the availability of volatiles and organics in the planetesimals, as these are thought to be heavily depleted in the warm, inner disk regions where habitable planets accrete. Organics and volatile content (interior or as a surface layer) change the optical properties of the dust grains, producing signatures that are detectable at optical and infrared wavelengths (e.g., Debes et al. 2008; Rodigas et al. 2014; Ballering et al. 2016). Recently discovered debris disks with gas content that may be recent or primordial provide an additional opportunity to explore volatile reservoirs in planetesimal belts (Dent et al. 2014; Kóspál et al. 2013; Moór et al. 2013)

4.2.3. Complementary Data

Exo-zodiacal disk studies will benefit from:

- 1) WFIRST imaging of debris disks;
- 2) ALMA observations of cold debris disks;
- 3) LBTI observations of the warm debris;
- 4) JWST observations of warm debris disks.

Observational Requirements

Sample size: Individual or small samples are useful.

Observations: Multi-wavelength optical/near-infrared/mid-infrared imaging, spectroscopy, and optical/near-infrared polarimetry. Large field of view ($\sim 10''$ or larger) may be needed to study disk morphology in nearby systems.

Questions to SAG15:

Input from WFIRST PS team on what debris disk science do they foresee.

5. Exoplanet Characterization

5.1. B1. How do rotational periods and obliquity vary with orbital elements and planet mass/type?

Contributors: Daniel Apai, Nicolas Cowan

A planet’s rotational state refers to both its obliquity and frequency, or equivalently period. Planetary rotation constrains the formation and angular momentum evolution of a planet, especially when comparing statistical samples of diverse planets. Moreover, the rotation of a given planet impacts its climate through diurnal forcing and its circulation through the Coriolis force, and contributes to magnetic field generation.

For example, Yang et al. (2014, 2013) showed that the rotation periods of temperate terrestrial planets changes the inner boundary of the habitable zone by a factor of two in insolation. Furthermore, planetary magnetic fields may be important shields against atmospheric loss. As these examples illustrate the rotational state of temperate terrestrial planets directly impacts their habitability.

5.1.1. *State of the Art to Measure Rotational Periods*

As of now little is known about the obliquity and rotational periods of non-synchronously rotating exoplanets. Rotational periods for planets and exoplanets have been determined through four different methods:

a) Phase Curve for Irradiated Planets: For some close-in synchronously rotating giant exoplanets the orbital/rotational phase modulation is detectable in the combined light of the star+planet system. The modulation also allowed longitudinal mapping the planets: For example, the dayside map of HD 189733b suggests that this hot Jupiter has zero obliquity (Majeau et al. 2012; de Wit et al. 2012). Although the eastward offset of the hotspot observed on most hot Jupiters (Knutson et al. 2007, 2009, 2012; Crossfield et al. 2010; Cowan et al. 2012b) is consistent with equatorial super-rotation on a synchronously-rotating planet (Showman & Guillot 2002), but also with slower winds on a non-synchronous planet (Rauscher & Kempton 2014). In fact, there is a complete degeneracy between the rotation of a gaseous exoplanet and its winds (Cowan & Agol 2011).

b) Period of the magnetic field’s rotation: The magnetic field is tracing the rotational periods of the planets’ *interiors*, which may be different from the latitude-averaged rotational periods measured in their upper atmospheres. In the Solar System Jupiter’s and Saturn’s

rotational periods are defined by the rotation of their inclined (w.r.t. spin axis) magnetic dipoles. For exoplanets, in exceptional cases, the manifestation of inclined magnetic dipoles may be detectable through time-varying auroral emission at UV/optical wavelengths or via modulated synchrotron emission in the radio. Recent detections of modulated radio emission from nearby brown dwarfs (e.g., Kao et al. 2016) suggests that very sensitive radio-wavelength observations of extrasolar giant planets may also be used in the future to establish their rotational periods.

c) Absorption line width measurements for directly imaged giant exoplanets (Beta Pic-toris b: Snellen et al. 2014). Similar studies for rotational line broadening have been carried out successfully for brown dwarfs (e.g., Reiners & Basri 2008). In order to convert the observed $v \sin i$ into a rotational period, one must know the planet’s radius and obliquity. This method is therefore well-suited for brown dwarfs and giant planets (which are all approximately the size of Jupiter), but could prove problematic for lower-mass directly-imaged planets of unknown radius. Furthermore, it is more applicable for systems where constraints exist on the planets’ obliquities.

d) Rotational photometric/spectroscopic modulations in hemisphere-integrated light for directly imaged exoplanets (Fig. 4, Zhou et al. 2016) and planetary-mass brown dwarfs (e.g., Biller et al. 2015, Leggett et al. 2016). Brown dwarfs (planetary mass and more massive), are good analogs for directly imaged exoplanets. These observations showed that low-level ($\sim 1\%$) rotational modulations in thermal emission are *very* common (Buenzli et al. 2014; Metchev et al. 2015), and can be used to measure or constrain rotational periods and to study cloud properties (e.g., Artigau et al. 2009; Radigan et al. 2012; Apai et al. 2013). Similarly, reflected-light observations of Solar System giant planets have also been used to demonstrate that rotational periods and their cloud covers can be characterized (e.g., Jupiter: Karalidi et al. 2015; Neptune: Simon et al. 2016).

Techniques may be both applicable for exoplanets directly imaged with next-generation space telescopes. While method *b* requires high spectral resolution and provides Doppler information, method *c* requires only high signal-to-noise time-resolved photometry and not strongly wavelength-dependent.

5.1.2. Science Cases

Habitable Planets (Earth-sized and Super-Earths): Rotation rates are an important parameter for climate and atmospheric circulation models of habitable planets: they constrain diurnal temperature modulations, determine the strength of the Coriolis force, in-




Fig. 3.— Whitened power spectrum from 50-day-long Kepler monitoring of hemisphere-integrated reflected light Neptune, with the most significant peak corresponding to the rotation period. Numbers above some peaks indicate the latitudes on Neptune corresponding to that rotation period based on the zonal velocities. From Simon et al. (2016).

fluence current and past magnetic field strengths and geometry, and indirectly constrain the atmospheric loss that may have occurred on these planets. Comparative studies of dynamo-generated magnetic energy densities in Solar System planets, the Sun, and rapidly-rotating low-mass stars show a correlation between the magnetic field strengths and the density and bolometric flux of the objects (see Fig. 5, Christensen et al. e.g., 2009). These studies argue for a scaling relation, based on Ohmic dissipation, where the field strength is only weakly sensitive to rotation rate, but the rotational rate fundamentally impacts the magnetic field geometry (bipolar vs. multi-polar, Christensen 2010). Furthermore, rotational rates also carry information about the accretion history of the planets and, in particular, about the size distribution of the planetary building blocks (e.g., Schlichting & Sari 2007).

In addition, the obliquity of habitable planets also has a major impact on the seasonal and diurnal temperature variations and on their climate in general. Obliquity is much more difficult to determine than the directly observable rotational rate. However, simulated observations demonstrate that it is possible to determine this quantity from high signal-to-noise reflected light lightcurves obtained at multiple orbital phases.

Considerable effort was put into exploring time-revolved observations of Earth, as exoplanet analog. Researchers have used simulated disk-integrated brightness variations of Earth to demonstrate that its rotational period can be estimated, even in the presence of time-varying clouds (Pallé et al. 2008; Oakley & Cash 2009). Likewise, such observations




Fig. 4.— Rotation periods provide insights into the properties and formation of planets. A comparison of Solar System planets, directly imaged exoplanets, and brown dwarfs reveals a characteristic mass-dependent rotation rate for massive planets. The arrows shows the expected spin-up due to gravitational contraction. From Zhou et al. (2016).

spanning multiple orbital phases constrain obliquity (Kawahara & Fujii 2010, 2011; Fujii & Kawahara 2012; Schwartz et al. 2016; Kawahara 2016). Schwartz et al. (2016) showed that although both latitudinal and longitudinal heterogeneities contribute to the obliquity signal, the latter contains more information. In principle, the amplitude modulation of rotational variations at only three orbital phases uniquely identifies a planet’s obliquity vector (the obliquity and its orientation with respect to the observer’s line of sight). Taking the complementary *frequency modulation* approach, Kawahara (2016) showed that modest signal-to-noise observations spanning most of a planet’s orbit could also constrain a planet’s obliquity, even if one is agnostic of the planet’s albedo map.

Atmosphereless Rocky Planets: Rocky planets with very thin or no atmosphere may exist (analogous to a “super-Mars” or a “dry Earth”, an Earth-like planet that lost its atmosphere and water). Such planets may form as a result of extensive atmospheric loss due to evaporation (Hot super-Mars), stellar wind stripping, or impact stripping (e.g., Schlichting et al. 2015). At pressures lower than water’s triple point (6 mbar) liquid water is not stable, even if the planet is otherwise Earth-sized and it is inside the habitable zone. The ability to measure rotational periods for these planets may provide important insights into the mechanism that led to the complete atmospheric loss. Atmosphereless rocky planets are suitable for direct measurements of their rotational periods as prominent albedo features

at the rocky surface will introduce photometric rotational modulations.

Fujii et al. (2014) used albedo-map generated lightcurves and, where available, observed photometric variations to explore the geological features detectable on diverse Solar System bodies with minor or no atmospheres (Moon, Mercury, the Galilean moons, and Mars). The study included the evaluation of the light curves and the features that are detectable at wavelengths ranging from UV through visible to near-infrared wavelengths, and also explored the accuracy required to determine the orbital periods of these bodies. Figure 6 provides an example for the wavelength-dependence of the rotational variability amplitudes in different bodies.

Gas and Ice Giant Exoplanets: The rotational periods of gas/ice giants may also be useful for constraining their formation and evolution (Tremaine 1991) and important for understanding their atmospheric circulation. Non-axisymmetrically distributed condensate clouds and hazes (photochemical or other origin) will introduce rotational modulations, both in reflected and in thermal emission (e.g., Simon et al. 2016). In addition, polarimetric modulations introduced by light scattering on heterogeneously distributed dust/haze grains may also be detectable. Currently, rotational rate estimates exist for close-in exoplanets (assumed to be equal to their orbital periods) and a few measurements exist for directly imaged exoplanets and planetary-mass brown dwarfs. The rotational angular momenta of close-in exoplanets (i.e., synchronously rotating) is reset by tidal interactions and no longer carries information on the intrinsic angular momenta of the objects. In contrast, the angular momenta of non-synchronously rotating exoplanets (such as those probed via direct imaging) carry information about their formation and angular momentum evolution. Photometric modulations have been measured in two near-infrared filters for the $\sim 4\text{--}6 M_{Jup}$ exoplanet 2M1207b (Zhou et al. 2016) and led to a rotational period measurement of $10.7_{-0.6}^{+1.2}$ h. CO absorption line rotational broadening measurements for the $10\text{--}13 M_{Jup}$ planet β Pictoris suggests a $v \sin i = 15$ km/s, which – assuming an equatorial viewing geometry, age, and mass – suggests a very similar rotational period. Similarly to these young exoplanets, photometric variations were used to measure the rotational periods of unbound young planetary mass-objects (Biller et al. 2015; Leggett et al. 2016) and very low-mass brown dwarfs (e.g., Scholz et al. 2015). The picture emerging – based on the very limited data – suggests that super-jupiter exoplanets and low-mass brown dwarfs start with similar angular momenta and during their evolution (cooling and contraction) their rotation rate increases, converging to the extrapolation of the Solar System mass-period relationship (see Figure 4).

A direct imaging mission capable of obtaining moderately high signal-to-noise ratio photometry of giant exoplanets can study possible trends between planet mass, semi-major axis, and rotational period.

Obliquity for gas giants: For gas giants (with well-constrained radius) combining the rotational period determined from rotational modulations with radial velocity information (line broadening due to rotation) allows constraining or deriving the inclination of the planet (e.g., Allers et al. 2016). Amplitude and frequency modulation of reflected light rotational variations (Schwartz & Cowan 2015; Kawahara 2016), Fourier spectrum or polarimetry of thermal emission (de Kok et al. 2011; Cowan et al. 2013).

A Note on Hazy Atmospheres: Planets with thick haze layers may pose a challenge for rotational signals using methods c and d (depending on the wavelengths of observations and the origins of molecular absorption or cloud features studied). Because haze particles *by definition* are small ($\sim 0.01 \mu\text{m}$), their residence time in the atmosphere will be much longer than the rotational period ($t_{res} \gg P$), which will result in featureless haze layers. As haze particles can be generated at smaller pressure levels and will settle down much slower than larger particles produced by condensation, the featureless haze layers *if optically thick* will mask any heterogeneous condensate cloud structure as well as any surface structures. Similarly, optically thick haze layers may cover or weaken the rotationally broadened line profiles in the atmospheres, also limiting the use of Doppler techniques. Therefore, planets enshrouded in thick haze layers are not well suited for rotational studies.

5.1.3. *The Science Value of Independently Measured Planet Masses and Radii*

Planetary Radius: For methods that measure rotational velocity the knowledge of planetary radius and obliquity is required to convert rotational broadening into a rotational period. However, if the goal is to determine the Coriolis forces, then rotational broadening is sufficient. For the photometric methods that produce a period estimate, on the other hand, the diurnal forcing pops out for free, while the Coriolis forces again require the planetary period. In general rotational information is most useful when combined with radius estimates. No complementary observations are required for science results from rotational period measurements, but observations constraining the planetary orbits may be combined with the obliquity and rotational period to constrain the formation history of low-mass planets.

Giant Planets: Radii for mature giant planets will be close to one Jupiter radius, but masses may vary by an order of magnitude. Masses may be derived from atmospheric modeling that includes a fit for surface gravity.

Radii and masses of rocky planets vary *more* than those of giant planets: mass may vary by a factor of ~ 20 (from Mars to super-Earths): while rotational periods alone will be important and useful for atmospheric circulation models, mass and/or radius measurements

would yield important additional science: mass measurements would allow exploring trends between formation mechanisms and angular momentum; and radius estimates (even from mass-radius relationships) would allow calculating Coriolis forces from rotational periods, significantly constraining the atmospheric circulation models.

Planet mass measurements from radial velocity or astrometry, or gravitational interactions between the planets, can be combined with rotational periods to determine the angular momenta of the giant planets, which may be useful for constraining their accretion history.

The periodicity in photometric variations is a direct measure of the rotational period, i.e., rotational period measurements do not require mass measurements. However, verifying the predicted trend between angular momentum, orbital period, mass (which potentially constrains the formation history) requires mass and radius measurements or uncertainties.

Observational Requirements (draft)

Sample size: Individual to medium/large

Observations: Very high spectral resolution; OR multi-epoch photometry.

Obliquity: photometry over complete rotational phase at three distinct orbital phases.

Pending Comments/Questions:

Kasper: ELT HCI + HRS will allow us to study rotation periods, sample will be small
- not sure about statistical relevance




Fig. 5.— The comparison between Earth, Jupiter, and stars shows that the magnetic energy density (in the dynamo) strongly correlates with a function of density and bolometric flux (here both in units of J m^{-3}). The bar lengths show estimated uncertainty rather than formal error (Supplementary Information). The stellar field is enlarged in the inset. Brown and grey ellipses indicate predicted locations of a brown dwarf with 1,500 K surface temperature and an extrasolar planet with seven Jupiter masses, respectively. From Christensen et al. (2009).




Fig. 6.— Albedo and its variations as a function of wavelengths for Solar System bodies with minor or no atmosphere. From Fujii et al. (2014).

Property	Optimal Wavelengths Range	Expected Amplitude	Acceptable Wavelengths Range	Expected Amplitude	Baseline or Num. Observations	Re
<i>Rotational Period</i>						
Earth/Super-Earth					3-30 h (1 Period)	
Super-Mars	0.9	25%	0.5-10	10-35%	3-30 h (1 Period)	7
Ice/Gas Giant	5	15%	0.3-5.0	3%	3-20 h (1 Period)	8
<i>Obliquity</i>						
Earth/Super-Earth						
Super-Mars	0.9	25%	0.5-1.0	10-35%	3 x 1 Period	7
Ice/Gas Giant	5	15%	0.3-5.0	3%	3 x1 Period	8

Table 1: My caption

5.2. B2: Which rocky planets have liquid water on their surfaces? Which planets have continents and oceans?

Relevance: Water is not a biosignature itself, but the presence of liquid water is required for life as we know it. Liquid water is not the only factor required for planetary habitability, but it is arguably the most important one. Thus, liquid water is a *habitability signature*. Establishing which habitable zone planets have liquid water on their surfaces provides an important context for EXOPAG SAG16, which focuses on biosignatures, but will rely on SAG15 for habitability signatures and characterization of habitable planets.

Our understanding of the distribution of water in exo-earths is very incomplete: Currently, water detections (direct and indirect) in extrasolar systems are limited to protoplanetary disks (e.g., Carr & Najita 2008; Salyk et al. 2008), hot jupiters and hot neptunes (e.g., Fraine et al. 2014), and in disks around white dwarfs fed by tidally disrupted minor bodies (Farihi 2016); however, no direct or indirect observations exist of water in extrasolar habitable zone Earth-like planets or even in super-Earths.

In the following we will discuss two different pathways for identifying liquid water on Earth-like habitable zone planets: 1) via the detection of oceans; and 2) via the inference of water vapor close to saturation pressure in their atmospheres.

5.2.1. Detecting Oceans

The traditional habitable zone (HZ) is defined in terms of surface liquid water (Kasting et al. 1993). Three distinct methods have been proposed to search for large bodies of liquids (oceans) on the surface of a planet:

Rotational color variability Oceans are darker and have different colors than other surface types on Earth, so the time variations in the color of a spatially unresolved planet can reveal the presence of liquid water oceans. This method relies on the existence of longitudinal heterogeneities in the planet’s surface composition (e.g., Ford et al. (2001); Cowan et al. (2009); Fujii et al. (2010); Kawahara & Fujii (2011,?)).

Polarization (e.g. Zugger et al. 2010, 2011). Oceans are smoother than other surface types (typically solids) and thus polarize light. For idealized scenarios, the phase variations in polarization are significant, but the same authors found that, in practice, the effect of oceans is masked by Rayleigh scattering, clouds, and aerosols. Observations of polarized Earthshine, however, imply that rotational variations in polarized intensity may be useful in detecting oceans (Sterzik et al. 2012).

Specular reflection (Williams & Gaidos 2008; Robinson et al. 2010). Oceans are also able to specularly reflect light, especially at crescent phases. The signal-to-noise requirements for phase variations are not as stringent as for rotational variations since the integration times can be much longer: weeks instead of hours. However, Robinson et al. (2010) showed that clouds not only mask underlying surfaces, but forward scattering by clouds mimics the glint signal at crescent phases, while atmospheric absorption and Rayleigh scattering mask the glint signature. They proposed using near-infrared opacity windows to search for glint, but this would only be possible if the effects of clouds could be accurately modeled for exoplanets. However, Cowan et al. (2012a) showed that crescent phases statistically probe the least-illuminated and hence coldest regions of a planet, insofar as these planets have ice and snow in their coldest latitudes, then this latitude albedo effect acts as false positive for ocean glint.

Although the faces of extrasolar planets will not be spatially resolved in the foreseeable future, their rotational and orbital motions produce detectable changes in color and brightness. Ford et al. (2001) used simulations of Earth to show that the changing colors of its disk-integrated reflected light encode information about continents, oceans, and clouds. The inverse problem – inferring the surface geography of a planet based on time-resolved photometry – is much more daunting than the forward problem.

Much progress has been made on the *exo-cartography* inverse problem since the seminal work of Ford et al. (2001). The rotational color variations of a planet can be used to infer the number, reflectance spectra, surface area, and longitudinal locations of major surface types (Fujii et al. 2010, 2011; Cowan et al. 2009, 2011; Cowan & Strait 2013). Meanwhile, the rotational and orbital color variations of an unresolved planet can be analyzed to create a 2-dimensional multi-color map – equivalently a 2D map of known surfaces (Fujii et al. 2010; Kawahara & Fujii 2011, 2010; Fujii & Kawahara 2012).

5.2.2. *Inferring Liquid Water*

Additional methods may be used to deduce the probable presence of liquid water on the surface of a potentially habitable planet without directly or indirectly detecting an ocean:

- 1) *Test whether partial pressure of water vapor reaches saturation, and/or*
- 2) *Identify clouds made of liquid water droplets (and not water ice).*

Analysis of simulated exo-earth observations was used to demonstrate that rotational phase mapping (time-resolved observations of hemisphere-integrated reflected light from the planet) can reveal the types and distribution of surfaces. Equipped with additional data on

the color/spectra of the features and the physical conditions on the planetary surface may be used to identify surface features as oceans and continents.

Add more description here.

Observational Requirements (draft)

Sample size:

Observations:

1. **Orbital semi-major axis** of a planet is critical as it may determine the presence of liquid water on the surface. How many visits per system are needed by a direct imaging mission to determine an accurate orbital distance?

2. **Presence of Greenhouses gases** and water vapor in the atmosphere: CO₂ and H₂O have strong features in the near-IR.

Comments on B2):

5.3. B3. What are the origins and composition of clouds and hazes in ice/gas giants and how do these vary with system parameters?

All Solar System planets with an atmosphere also harbor cloud and/or haze layers. Clouds and hazes influence the pressure-temperature structure of the atmosphere, its emission and transmission spectra, as well as its albedo. Particles or droplets that make up clouds primarily form through condensation and grow via further condensation and/or particle collisions. With grain sizes that may range from a micron to millimeter cloud particles/droplets have short settling time and are typical at higher atmospheric pressures (~ 1 bar). Based on different extrapolations of clouds observed on Earth and on other Solar System clouds, a range of cloud models have been proposed for giant exoplanets and brown dwarfs (for a review and comparison, see Helling et al. 2008). Haze particles (typically $< 0.1 \mu\text{m}$ in size) often form via photochemistry-driven (e.g., Titan) or charged-particles-driven chemical reactions in the upper atmospheres (< 1 bar); with long residence times these particles often introduce large optical depths to upper atmospheres. From an observational perspective clouds and hazes may also be used as tracers of atmospheric dynamics (circulation, mixing, turbulence). Presence of haze or cloud layers may also mask the presence of specific atmospheric absorbers even if present at large abundances at pressures higher than the particle layer.

Current Knowledge: Condensate clouds have been observed in brown dwarfs and in hot Jupiters, over a very broad range of temperatures and pressures. High-altitude haze layers have been observed for transiting planets ranging from hot Jupiters to super-Earths and possibly for Earth-sized planets, as well as for brown dwarfs. In the following we briefly summarize the key aspects of condensate clouds and haze layers and refer the reader to one of the many excellent in-depth reviews of the field for more complete summary.

Condensate Clouds: As the atmospheres of exoplanets encompass a very broad temperature range (~ 200 to $2,000$ K) these atmospheres are expected to harbor a large variety of condensates. For solar compositions the most important condensates include Ca-Ti-oxides, silicates, metallic iron, sulfides, CsCl and KCl, H_2O , NH_4HS , NH_3 (e.g., Lodders & Fegley 2002, for a recent review see Marley & Robinson e.g., 2015). Most of our current knowledge on cloud properties and compositions come from studies of Solar System planets (most importantly, Earth and Jupiter) and from the abundant samples of brown dwarfs. Water vapor and water ice clouds in Earth can be studied in-situ and via remote sensing; models developed to explain their behavior and properties are often used as a starting point for models of extrasolar clouds (Ackerman & Marley 2001), although it is likely that in some exoplanet and brown dwarf atmospheres cloud formation and properties may be set by different processes (for a review of different cloud models see, e.g., Helling et al. 2008).

With over $\sim 3,000$ brown dwarfs known these objects provide a easy-to-study analogs of extrasolar giant planet atmospheres. Temperatures of known brown dwarfs range from ~ 250 K (below freezing point!) to above 2,300 K; an increasing number of known brown dwarfs have very low gravities and masses of only a few M_{Jup} , enabling the definition of samples essential for comparative parameter studies.

Comparative studies of brown dwarfs reveal the presence of silicate cloud layers through prominent infrared color-magnitude changes that occur through the M–L–T–Y spectral type sequence. The sequence itself is primarily set by the presence and absence of prominent gas-phase absorbers and not directly by the presence/absence of clouds (e.g., Burgasser et al. 2006; Cushing et al. 2006; Kirkpatrick et al. 2012); however, there is a strong correlation between the spectral type and colors of a given object (e.g., (Burgasser et al. 2002; Burrows et al. 2006; Saumon & Marley 2008; Dupuy & Liu 2012)). The general and oversimplified picture that emerged suggests that while the hottest (M-type) brown dwarfs are cloud free, with temperatures sinking below $\sim 1,800$ K the atmospheres of L-type brown dwarfs is characterized by thick silicate clouds (resulting in red near-infrared colors); at lower temperatures ($T < 1,300$ K) a transition to silicate cloud-free atmospheres is envisioned. Correspondingly, cool T-type brown dwarfs have blue near-infrared colors, consistent with the lack of thick clouds in their upper atmospheres (see Figure 8). At even lower temperatures, within the Y spectral type, less refractory and less abundant species, including water ice, are expected to condense out and form clouds (e.g., Morley et al. 2014).




Fig. 7.— Condensate clouds predicted for the upper atmospheres of planets of different temperature. By D. Apai, after Lodders (2003).

Although there is ample evidence supporting the overall picture described above, it is also clear that the above picture fails to capture the real complexity of cloud properties and




Fig. 8.— Condensate clouds have a fundamental impact on the positions of brown dwarfs and directly imaged exoplanets on the near-infrared color-magnitude diagram. Along the L-type sequence (red) silicate clouds in the upper atmosphere become thicker. The cooler T-dwarfs are bluer because the silicate clouds are below the visible upper atmosphere. Figure from Wagner et al. (2016), which is in part based on the parallax database by Dupuy & Liu (2012).

atmospheric chemistry in brown dwarfs. Outstanding questions include the large dispersion in color along the L–T sequence (e.g., Burgasser et al. 2008); the unusually red colors of many of the very young brown dwarfs (and a few intermediate-age ones), a likely sign of unusually dusty upper atmospheres (e.g., Allers & Liu 2013; Liu et al. 2013). Furthermore, the first detections of water ice clouds has been reported in a Y-dwarf with an effective temperature of only ~ 250 K (Faherty et al. 2014; Skemer et al. 2016), enlarging the temperature range over which cloud models can be tested.

Recently, time-resolved high-precision observations (photometric and spectroscopic light curves) enabled the comparative studies of different cloud layers within the same objects, breaking the degeneracy between the effects of the multiple atmospheric parameters that may vary between any two brown dwarf (age, composition, temperature, surface gravity, vertical mixing, cloud structure). Space-based (HST and Spitzer) studies with sub-percent photo-




Fig. 9.— Gemini/GNIRS spectrum of the ~ 250 K Y-dwarfs WISE0855 shows a series of absorption features attributable to water vapor, muted by clouds (likely water ice). From Skemer et al. (2016).

metric precision found that most, if not all, brown dwarfs have heterogeneous (patchy) cloud cover (Buenzli et al. 2014; Metchev et al. 2015); ground-based surveys found that the highest amplitude brown dwarfs are at the L/T transition Radigan et al. (2014). Time-resolved spectroscopy of L/T transition dwarfs showed that the spectroscopic variations emerge from the atmospheres characterized by a mixture of warm thin cloud / cooler thick cloud patches, and *not* by clouds and cloud holes (Apai et al. 2013; Buenzli et al. 2015). Simultaneous HST (1.1-1.7 μm) and HST–Spitzer (1.1-1.7 μm and [3.6] or [4.5]) observations of clouds in L, L/T, and T-type brown dwarfs revealed pressure-dependent (vertical) structures with characteristic patterns for objects of different spectral types (Buenzli et al. 2012; Apai et al. 2013; Yang et al. 2015). Most recently, planetary-mass brown dwarfs and companions have also been accessible to rotational modulation studies, providing an opportunity to explore cloud properties as a function of gravity (Biller et al. 2015; Zhou et al. 2016; Leggett et al. 2016).

Directly imaged exoplanets and planetary-mass companions cover a spectral type range from early L to mid-T. These objects differ from old, high-gravity brown dwarfs both in the fine structure of their spectra (Barman et al. 2011,?; Skemer et al. 2012) and, often, in their broad-band colors (see, e.g., Fig. 8), but show some strong similarities to some young brown dwarfs (e.g. Allers & Liu 2013; Faherty et al. 2013, 2016). From the small sample of directly imaged exoplanets it appears that early L-type exoplanets have colors similar to brown dwarfs with matching spectral types (), late L and L/T-type exoplanets are often much redder and fainter than brown dwarfs with matching spectral types (e.g., Chauvin

et al. 2005; Marois et al. 2008), but the coolest T-type exoplanets appear to have colors consistent with those of T-type brown dwarfs (Macintosh et al. 2015; Wagner et al. 2016). This pattern, if verified, would argue for a difference in cloud properties (most significant in late-L and L/T transition objects) between the higher gravity brown dwarfs and the low-gravity exoplanets.

Clouds have also been studied in hot jupiters via transmission and emission spectroscopy, spectral phase mapping, and in reflected light. Observations from the Kepler space telescope (dominated by reflected light) argued for a large-amplitude, heterogeneous silicate cloud cover (e.g., Demory et al. 2013) that avoids the cold trap in the night side of the planet. (Fortney et al. 2008) proposed that the presence/absence of silicate clouds in hot jupiters should follow the general sequence observed in brown dwarfs. Although optical-near infrared HST transmission spectra argued for the presence of cloud decks in some hot jupiters (Gibson et al. e.g., 2013; Sing et al. e.g., 2015), no clear trend (in terms of presence/absence of clouds) emerged from a homogeneous survey of hot jupiters (Sing et al. 2016). It is likely that the presence of silicate clouds in the regions probed by transmission (terminator) and emission spectroscopy (dayside) strongly depends on the day-night temperature difference (e.g., Rauscher & Menou 2013), atmospheric circulation (see Question C1, and e.g., Showman et al. 2009, 2015), and the importance of potential cold traps (Parmentier et al. 2013).

Hazes – with particles less than $0.1 \mu\text{m}$ – have been argued for in a few objects where the lack of near-infrared absorption features (commonly water) necessitates that the absorption features are muted by high-altitude particles (unless the upper atmospheres of some transiting planets are extremely dry, see Madhusudhan et al. 2014). Such strong reduction in water absorption features was seen in the warm sub-neptune GJ1214b (Kreidberg et al. 2014) and atmospheric models argued for the presence of very small particles at low pressures (~ 1 mbar), consistent with photochemical hazes but not with condensate clouds (Morley et al. 2015). In an ongoing study, the transmission spectra of moderate earth-sized planets around a very low-mass red dwarf also appear to be flat, perhaps also influenced by small particles lofted to low pressures (de Wit et al. 2016, in prep.). The detection of hazes (based on a very similar water absorption-based evidence) argues for the presence of some haze in L-type brown dwarfs, in spite of the lack of a host star, which argues for non-photochemical haze production (Yang et al. 2015), possibly driven by charged particles accelerated by the brown dwarf’s magnetic field.

Solar System Gas Giants as exoplanet analogs observations: Overlapping Kepler photometry and Hubble Space Telescope images of Neptune have shown complex time-varying signal whose frequency analysis revealed not only the fundamental rotation rate, but also the level of differential rotation of major mid-latitude cloud features Simon et al. (2016). Quasi-

continuous 20-hour-long two-band optical imaging of Jupiter with the Hubble Space Telescope provided simultaneous high-precision photometry and high-fidelity and high-resolution images (Karalidi et al. 2015). These authors showed that MCMC-based lightcurve modeling can correctly retrieve the position, size, and surface brightness of the dominant features in the lightcurve, such as the Great Red Spot, even from a single rotation.

Sub-questions

The study of extrasolar cloud layers is novel and we may not be in the position yet to identify the right set of key questions to ask. Nevertheless, the following list attempts to capture the most important uncertainties of our current models of clouds.

- What are vertical structures of single and multi-layer clouds formed from different condensates?
- What are the grain size distributions and compositions (single-species or compound grains) in the clouds?
- Under which conditions do photochemically- and charged particle-driven haze layers form? How complex can chemistry get in haze layers?
- How do condensate clouds form and evolve as a function of fundamental atmospheric parameters?

Observational Requirements (draft)

Sample size:

Observations:

Questions to SAG15:

- 1) How challenging are the very different methods described here relative to each other?
- 2) How are they best carried out?

Kasper: ELT: A Jupiter analogue (1Rj, 5AU) will just be observable ($c \sim 1e-9$). Neptune / Uranus analogues cannot be seen with an ELT.

5.3.1. Science Value of Independently Measured Planet Masses and Radii

6. Exoplanetary Processes

6.1. C1. What processes/properties set the modes of atmospheric circulation and heat transport in exoplanets and how do these vary with system parameters?

Authors: Daniel Apai, Nick Cowan, Ravi Kopparapu

Atmospheric circulation plays a key role in redistribution energy in exoplanet atmospheres. Depending on typical wind speeds, rotational velocity, insolation, latent heat released during condensation, and other system parameters different atmospheric circulation regimes are expected on planets that can be studied with direct imaging missions. For potentially habitable exoplanets atmospheric circulation will determine the day-night heat differential and the equator-pole temperature difference. Understanding the presence and size of Hadley cells can also provide important insights into how water vapor (or other condensibles) may be distributed in habitable planets.

Understanding atmospheric circulation in habitable exoplanets is an important component in establishing a correct climate model for them. As of now, atmospheric circulation has been probed in the Solar System planets and a small sample of brown dwarfs, hot jupiters and lower-mass exoplanets (see Figs 11 and 12, Yang et al. 2013; Leconte et al. 2013; Abe et al. 2011; Wordsworth et al. 2011; Zhang & Showman 2014; Kopparapu et al. 2016; Kataria et al. 2014; Wolf & Toon 2014). The nature of the atmospheric dynamics depends upon how thick an atmosphere the planet has, the rotation rate of a planet, the distance of the planet from the star and several other factors. A more comprehensive study of different atmospheric circulation regimes of exoplanets remain unexplored. Using time-resolved observations and rotational phase mapping techniques atmospheric circulation may be constrained.

Questions to SAG15:

To what level can the atmospheric circulation be constrained for different types of planets?

What hypotheses / toy circulation models should be tested for gas giants?

What hypotheses / toy circulation models should be tested for habitable super-earths / earths?

What data type and cadence is required or best suited for characterizing circulation?

Observational Requirements (draft)

Sample size:




Fig. 10.— Depending on the relative importance of rotational speed, wind speed, and vertical heat transport, simple models predict two different regimes of circulation for giant planets: vortex-dominated (left) and jet-dominated (right). From Zhang & Showman (2014).




Fig. 11.— Moist/water-rich atmosphere simulations from Wolf & Toon(2014). The four panels indicate the amount of cloud fraction on a planet at different insolation (or alternately how close to an inner edge of the HZ a planet is located). From left to right, the solar insolation varies S_0 (current Earth insolation), 10% of S_0 , 12.5% S_0 and 21% of S_0 . This is for an Earth-size planet around a Sun-like star.




Fig. 12.— Temperature and horizontal wind vectors at the surface, 0.5 bar, and 0.1 bar levels for an Earth-mass planet in a slow-rotating regime near the inner edge of the habitable zone around a K-dwarf. Slowly rotating planets develop subs-stellar clouds that increase the albedo of the planet. Inflow along the equator and from the poles into the substellar point at the center is also shown. From Kopparapu et al. (2016).

Observations:

6.2. C2. What are the key evolutionary pathways for rocky planets and what first-order processes dominate these?

Contributors: Nick Cowan, Daniel Apai

The two earth-sized rocky planets in the Solar System, Earth and Venus, likely have started with very similar initial mass, orbit, and composition, but their evolutionary paths have strongly diverged. Mars, although substantially different in its mass and orbit, has again followed a different evolutionary trajectory, even though it is thought that surface conditions at the early Mars, at least temporarily or episodically, may have resembled the early Earth. With the large number of rocky planets that may be observable with a capable future direct imaging mission the range of evolutionary histories of planets could be explored.

The questions naturally emerge: What key evolutionary pathways exist for rocky planets and what factors determine which of these pathways a given planet will follow?

Attractors and Divergence in the Phase Space of Rocky Planet Evolution: It is reasonable to describe the momentary state of a given rocky planets with a set of n fundamental parameters and explore the evolution of the planet in this n -dimensional phase space. Each planet's history and future evolution is thought of as a trajectory. Fundamental parameters could include, but not be limited to, planet mass, radius, atmospheric pressure scale height, orbital parameters, atmospheric composition, magnetic field strength, etc. Which trajectory a planet follows will depend not only in its momentary location in the phase space, but also by the effect of a set of feedback loops (both positive and negative) as well as on a few environmental variables (e.g., stellar luminosity and incident optical and UV flux).

When describing planet evolution in such a manner, several obvious questions are identified: 1) *How sensitive are the trajectories to initial parameters and/or perturbations to the system?* 2) *What is the importance of a planet's past, e.g., which volumes of the phase space are uni-directional (e.g., irreversible water loss)?* 3) *Are there preferred evolutionary end-states ('attractors') or is the surface defined by coeval planets smooth?* 4) *What is the importance of quasi-monotonic evolution driven by a small number processes vs. random walk driven by a multitude of competing processes?*

Exploring the past history and current state of rocky planets allows the system-level study of rocky planet evolution and will be essential for understanding the occurrence rate of truly earth-like planets and to place the physical processes that drive planet evolution on

Earth to the broader context of exo-earths.

6.2.1. *Science Value of Independently Measured Planet Masses and Radii*

Is the science goal achievable without precise mass measurements? Yes, a medium or large sample of rocky exoplanets for which most of the other key parameters are known would likely suffice to establish the topology of the phase space.

Would the science goal benefit greatly from precise mass measurements? Yes, precise mass measurements would significantly contribute to the understanding of the planets' properties. In case the phase space is highly complex and its projection to a lower-dimensional (observed) phase space does not allow the identification of the key processes that drive the evolution, expanding the projected phase space by a new dimension (mass) may break the degeneracy between different processes that lead to similar evolutionary outcomes.

Observational Requirements (draft)

Sample size: medium to large (>15)

Observations: characterization of the planets: atmosphere pressure and composition, orbital parameters, bulk composition, surface temperature estimate, stellar parameters and past evolution;

Comments: Solicit input from: Robert Boschia, Valerio Lucarinia, Salvatore Pascalea (2013); William Moore, Adrian Lenardic, Dorian Abbot, Dan Fabrycky

To explore: Toy model for testing hypothesis of smooth distribution vs. attractors

6.3. C3. What types/which planets have active geological activity, interior processes, and/or continent-forming/resurfacing processes?

Contributors: Stephen Kahn, Daniel Apai, Nick Cowan

Planetary interior processes and geological activity play an important role in coupling Earth's atmosphere to its crust and providing a long-term stabilizer for Earth's climate. The source of Earth's atmosphere and volatiles are mostly products of outgassing after the loss of the primary atmosphere. Developing reliable climate models to determine the habitability of potentially habitable planets will likely require assumptions on the geological activity and the level of coupling between the planet's crust and atmosphere. Interior processes are obviously very difficult to probe via low signal-to-noise and spatially unresolved remote sensing.

The influence of volcanism on planetary climate is most clearly understood for the case of Earth. On geologic timescales, continental crust production participates in the stabilization of the Earth’s climate through its role in carbonate weathering feedback. Chemical weathering of silicate minerals on land in the presence of water causes the slow removal of CO_2 from the atmosphere, which is eventually deposited on the ocean floor as carbonate compounds. Without the continual re-injection of new CO_2 by volcanoes, the atmospheric stock of CO_2 would be slowly depleted. However, the rate of CO_2 removal by silicate weathering is temperature dependent, so that in the presence of a steady source of volcanic CO_2 , weathering interacts with the greenhouse properties of CO_2 to produce a negative feedback on planetary temperature. This interaction, whereby warmer conditions lead to increased drawdown of CO_2 and a consequent weakening of the greenhouse effect (and vice versa), is believed to play an important role in stabilizing planetary temperatures in the presence of a main-sequence star which is increasing in luminosity over Ga timescales. It is because of this process that it has been argued that volcanism and geological activity are necessary conditions for sustained life on a planet.

Current Knowledge: Currently no methods are capable of probing geological activity in exoplanets and our knowledge is limited to a range of possible activity levels predicted by various extrapolations of the activity levels of rocky planets in the Solar System.

Studies of terrestrial climate and volcanism focus primarily on the effects of volcanism on surface temperature, which we are unlikely to be able to estimate for most exoplanets. However, volcanically forced anomalies in surface temperature are coupled to anomalies in emission temperature, which can be targeted for follow-up observations. Thus, if volcanism can be identified on an exoplanet it may represent the most promising method for estimation of climate sensitivity outside of the Solar System.

The distinctive effect of volcanic eruptions on the transmissivity of atmospheres is related to the force of their explosions. Typically, processes on Earth that produce aerosols in the atmosphere affect only the troposphere. Aerosols are quickly washed out of the troposphere by rain, and thus do not have a sustained impact on atmospheric transmissivity. Many small eruptions don’t reach the stratosphere, however the largest explosive volcanic material can, in contrast, inject SO_2 directly into the stratosphere, where it reacts to form sulphate aerosols.

Because the stratosphere is very dry, these aerosols can persist in the stratosphere for several years, until they are removed by the natural overturning circulation of the stratosphere (Robock et al. 2007). Stratospheric air rises in the tropics and then migrates towards the pole where it sinks. Because of this, aerosols from tropical eruptions typically persist in the stratosphere for about two years, while aerosols from high-latitude volcanism persist for

only one year (Robock et al. 2007; Tingley et al. 2014).

Previous work shows a link between exoplanet compositions and stellar compositions (e.g., Rogers & Seager 2010) such that stellar compositions can be used to approximate the composition of exoplanet interiors. Stars in exoplanetary systems show a wide variation in composition (Hinkel et al. 2014). In particular, some composition parameters with large variability such as Mg:Si ratios, are likely to have a first order effect on the minerals that compose exoplanetary interiors and thus the melting behavior, magma composition generated from these planetary mantles, and their volatile solubility. Certain compositional components, such as alkalis, have also been shown to greatly increase the H₂O solubility (e.g., Behrens & Zhang 2001; Larsen & Gardner 2004) in natural melts, and highlight the necessity of measuring volatile solubility behavior across a broad range of melt compositions. Magmatic volatile solubility is highly dependent on temperature, which also varies with mineralogy. For example, we hypothesize that planets with bulk mantle compositions that are dominated by pyroxene and feldspar type minerals (Mg:Si < 1) will have lower melting temperatures and thus higher H₂O solubility than planets dominated by olivine.

On Earth, in addition to the pressure- and compositional-dependence of volatile solubility in magmas, the explosivity of a given eruption is dependent on the overall volatile concentration (dominated by H₂O and CO₂), magma supply rate, vent geometry, and source pressure of the magma body (e.g., Wilson 1980; Papale & Polacci 1999; Mason et al. 2004). The most explosive eruptions on Earth tend to be those at convergent plate boundaries where there are abundant volatiles involved in magma genesis sourced from the subducting plate, and some types of intraplate volcanism where interactions with reservoirs of volatiles in the crust produce highly explosive caldera eruptions. In addition, flood basalts and other volumetrically large outpourings of magma common in a planets early history may be a significant source of atmospheric volatiles (Black et al. 2012). As such, the lack of tectonics on exoplanets does not preclude extreme volcanism that would likely produce detectable signatures.

O’Neill et al. (2013) provides a recent review of plate tectonics in Earth and Venus and its sensitivity to radioisotope abundances.

6.3.1. Geological Activity and Plate Tectonics on Extrasolar Rocky Planets

The extrapolation of models of planetary evolution and plate tectonics to extrasolar rocky planets is challenging. A particularly relevant question is how plate tectonics may operate in super-Earths: on one hand, the higher heat flux (due to their intrinsically higher

mass-to-surface ratio) should lead to stronger mantle convection (e.g., Valencia et al. 2007; van Heck & Tackley 2011). On the other hand, based on a visco-elastic models of mantle convection and crust formation, O’Neill & Lenardic (2007) find that increasing the planet’s radius (and mass) will decrease the ratio of driving-to-resistive forces (see Fig. 13), which reduces the likelihood of mobile plate tectonics in super-Earths and argues for the stagnant lid (or episodic tectonics) in these planets.

Furthermore, for a given planet models also suggest time-dependence and sensitivity to initial conditions: the thermal state of the post-magma ocean mantle is a key parameter that determines the subsequent evolution of the planet (possibly but not necessarily through i) hot stagnant-lid, ii) plate tectonics, then to iii) cold stagnant lid regime). Depending on the planet’s transition from the magma ocean stage different evolutionary paths are possible and there may only be a limited time available for Earth-like plate tectonics O’Neill et al. (2016).

6.3.2. *Observational Methods*

While major geological processes usually unfold on timescales not accessible to large-distance remote sensing, the *results* of these processes are detectable and, in some cases, may be unambiguously identifiable. For example, in the case of Earth the presence of multiple large land-masses and oceans (detectable via time-resolved observations, e.g., Cowan et al. 2009) reveals that a continent-forming process acts on timescales shorter than water-driven land erosion and provides a characteristic scale for the continental plates. Another Earth-based example is the accumulation of atmospheric absorbers characteristic volcanic outgassing (e.g., SO₂: Kaltenegger & Sasselov 2010). Other, non-Earth-like, planets may offer other detectable signatures of geological activity.

In the following we briefly discuss four representative possibilities:

- i) Continents and Oceans from Surface Maps
- ii) Atmospheric Absorbers from Volcanic Outgassing
- iii) Planetary-Scale Surface Mineralogy
- iv) Cloud formation as Tracer of Topography and Erosion

Continents and Oceans from Surface Maps: Simulated observations of Earth as an exoplanet demonstrate that with appropriate rotational- and orbital phase-resolved precision,




Fig. 13.— Convection as a function of stellar radius and Byerlee-style pressure-dependent yield stress. The models include internal heating, a constant friction coefficient, and gravity matching the planetary mass. Larger radius results in greater buoyancy forces, but also increased fault strength due to increased pressure. Thus planets with larger radii again tend to be in an episodic or stagnant regime, depending on the absolute yield stress. From O’Neill & Lenardic (2007).

multi-band photometric data can be used to identify the presence and one-dimensional and two-dimensional distribution of oceans and landmasses (see also Section 5.2.1, Cowan et al. e.g., 2009; Fujii et al. e.g., 2010; Fujii & Kawahara e.g., 2012). In a planet where large bodies of liquid water (an ocean) is present, a hydrological cycle is active, and land masses (continents) are detected, land erosion is arguably must occur; timescales for the erosion can be estimated based on terrestrial measurements. The existence of the continents demonstrates that the time-scale of continent-formation is comparable or faster than their erosion. Based on a simplified model of water cycling and continent formation Cowan & Abbot (2014) argues that continents and oceans may be common even among super-earths with high abundances of water. Such first-principle-based models may be combined with the scales of oceans and continents derived from observations to test whether active continent formation (e.g., plate tectonics) is required for a given planet.

Water Clouds as Tracer of Topography Similarly to the previous Pallé et al. (2008)

Planetary-scale Surface Geology Fujii et al. (2014) used albedo-map generated lightcurves and, where available, observed photometric variations to explore the geologic features detectable on diverse Solar System bodies with minor or no atmospheres (Moon, Mercury, the Galilean moons, and Mars). The study included the evaluation of the light curves and the features that are detectable at wavelengths ranging from UV through visible to near-infrared wavelengths, and also explored the accuracy required to determine the orbital periods of these bodies. Figure 6 provides an example for the wavelength-dependence of the rotational variability amplitudes in different bodies.

Amplitude variations at the level of 5–50% have been reported introduced by features of diverse nature (volcanism, space weathering, planetary weathering, impact excavation, tectonic deformation). In some cases data with the appropriate wavelength coverage can be used to identify some of these features or narrow down the possible origins.

Volcanic Outgassing

6.3.3. Complementary Datasets

We identify three complementary datasets that are critically important for modeling the interior and activity of extrasolar rocky planets:

- *Stellar abundances*: a proxy for the relative refractory elemental abundances that may be present in the planet; may be used to identify outlier systems in terms of elemental abundances

- *Composition of giant planets in the system:*
- *Stellar/system age:* to constrain the evolutionary state of the planets (heat flux and time available for volatile loss and resurfacing processes)
- *Mass and radius of the planet:* fundamental physical parameters with major impact on energy budget and force balances; and constrains the bulk composition

6.3.4. *Science Value of Independently Measured Planet Masses and Radii: Very High*

Exploring the planetary-scale geophysics of rocky planets will likely be among the most challenging aspects of characterizing extrasolar rocky planets. Yet, understanding the geophysics and interior activity of these planets may well turn out to be essential for correctly and robustly interpreting atmospheric biosignatures. The rocky planet’s mass is one of the most fundamental parameter that influences heat flux, pressure, and horizontal forces acting on the lithosphere. Given the sensitivity of plate tectonics models to planet mass, it is likely that determining the planet mass with a precision of $\sim 10\%$ is required for establishing a robust geophysical model.

7. Data Requirements

This section identifies the type and quality of data ideal or required to answer the individual science questions..

Appendix

A. SAG15 Charter

Future direct imaging missions may allow observations of flux density as a function of wavelength, polarization, time (orbital and rotational phases) for a broad variety of exoplanets ranging from rocky sub-earths through super-earths and neptunes to giant planets. With the daunting challenges to directly imaging exoplanets, most of the community's attention is currently focused on how to reach the goal of exploring habitable planets or, more specifically, how to search for biosignatures.

Arguably, however, most of the exoplanet science from direct imaging missions will not come from biosignature searches in habitable earth-like planets, but from the studies of a much larger number of planets outside the habitable zone or from planets within the habitable zone that do not display biosignatures. These two groups of planets will provide an essential context for interpreting detections of possible biosignatures in habitable zone earth-sized planets.

However, while many of the broader science goals of exoplanet characterization are recognized, there has been no systematic assessment of the following two questions:

- 1) What are the most important science questions in exoplanet characterization apart from biosignature searches?
- 2) What type of data (spectra, polarization, photometry) with what quality (resolution, signal-to-noise, cadence) is required to answer these science questions?

We propose to form SAG15 to identify the key questions in exoplanet characterization and determine what observational data obtainable from direct imaging missions is necessary and sufficient to answer these.

The report developed by this SAG will explore high-level science questions on exoplanets ranging from gas giant planets through ice giants to rocky and sub-earth planets, and – in temperatures – from cold (~ 200 K) to hot ($\sim 2,000$ K). For each question we will study and describe the type and quality of the data required to answer it.

For example, the SAG15 could evaluate what observational data (minimum sample size, spectral resolution, wavelength coverage, and signal-to-noise) is required to test that different formation pathways in giant planets lead to different abundances (e.g. C/O ratios). Or the SAG15 could evaluate what photometric accuracy, bands, and cadence is required to identify

continents and oceans in a habitable zone Earth-sized or a super-earth planet. As another example, the SAG15 could evaluate what reflected light data is required to constrain the fundamental parameters of planets, e.g. size (distinguishing earth-sized planets from super-earths), temperature (cold/warm/hot), composition (rocky, icy, gaseous), etc.

SAG15 will not attempt to evaluate exoplanet detectability or specific instrument or mission capabilities; instead, it will focus on evaluating the diagnostic power of different measurements on key exoplanet science questions, simply adopting resolution, signal-to-noise, cadence, wavelength coverage as parameters along which the diagnostic power of the data will be studied. Decoupling instrumental capabilities from science goals allows this community-based effort to explore the science goals for exoplanet characterization in an unbiased manner and in a depth beyond what is possible in a typical STDT.

We envision the SAG report to be important for multiple exoplanet sub-communities and specifically foresee the following uses: 1) Future STD teams will be able to easily connect observational requirements to missions to fundamental science goals; 2) By providing an overview of the key science questions on exoplanets and how they could be answered, it may motivate new, dedicated mission proposals; 3) By providing a single, unified source of requirements on exoplanet data in advance of the Decadal Survey, the science yield of various missions designs can be evaluated realistically, with the same set of assumptions.

Our goal is to carry out this SAG study by building on both the EXOPAG and NExSS communities.

We aim to complete a report by Spring 2017 and submit it to a refereed journal, although this timeline can be adjusted to maximize the impact of the SAG15 study for the ongoing and near- future STDTs and other mission planning processes.

Synergy with a potential future SAG proposed by Shawn Domagal-Goldman: While the SAG proposed here will include studies of habitable zone rocky planets, it will focus on planets without significant biological processes. A future SAG may be proposed by Shawn Domagal- Goldman to explore biosignatures; if such a SAG is proposed, we envision a close collaboration on these complementary, but distinct problems.

REFERENCES

- Abe, Y., Abe-Ouchi, A., Sleep, N. H., & Zahnle, K. J. 2011, *Astrobiology*, 11, 443
- Ackerman, A. S. & Marley, M. S. 2001, *ApJ*, 556, 872
- Allers, K. N., Gallimore, J. F., Liu, M. C., & Dupuy, T. J. 2016, *ApJ*, 819, 133
- Allers, K. N. & Liu, M. C. 2013, *ApJ*, 772, 79
- Apai, D., Radigan, J., Buenzli, E., et al. 2013, *ApJ*, 768, 121
- Apai, D., Schneider, G., Grady, C. A., et al. 2015, *ApJ*, 800, 136
- Artigau, É., Bouchard, S., Doyon, R., & Lafrenière, D. 2009, *ApJ*, 701, 1534
- Ballering, N. P., Su, K. Y. L., Rieke, G. H., & Gáspár, A. 2016, *ApJ*, 823, 108
- Barman, T. S., Macintosh, B., Konopacky, Q. M., & Marois, C. 2011, *ApJ*, 733, 65
- Behrens, H. & Zhang, Y. 2001, *Earth and Planetary Science Letters*, 192, 363
- Biller, B. A., Vos, J., Bonavita, M., et al. 2015, *ApJ*, 813, L23
- Black, B. A., Elkins-Tanton, L. T., Rowe, M. C., & Peate, I. U. 2012, *Earth and Planetary Science Letters*, 317, 363
- Boccaletti, A., Thalmann, C., Lagrange, A.-M., et al. 2015, *Nature*, 526, 230
- Buenzli, E., Apai, D., Morley, C. V., et al. 2012, *ApJ*, 760, L31
- Buenzli, E., Apai, D., Radigan, J., Reid, I. N., & Fplateau, D. 2014, *ApJ*, 782, 77
- Buenzli, E., Saumon, D., Marley, M. S., et al. 2015, *ApJ*, 798, 127
- Burgasser, A. J., Geballe, T. R., Leggett, S. K., Kirkpatrick, J. D., & Golimowski, D. A. 2006, *ApJ*, 637, 1067
- Burgasser, A. J.,Looper, D. L., Kirkpatrick, J. D., Cruz, K. L., & Swift, B. J. 2008, *ApJ*, 674, 451
- Burgasser, A. J., Marley, M. S., Ackerman, A. S., et al. 2002, *ApJ*, 571, L151
- Burrows, A., Sudarsky, D., & Hubeny, I. 2006, *ApJ*, 640, 1063
- Carr, J. S. & Najita, J. R. 2008, *Science*, 319, 1504

- Chauvin, G., Lagrange, A.-M., Dumas, C., et al. 2005, *A&A*, 438, L25
- Christensen, U. R. 2010, *Space Sci. Rev.*, 152, 565
- Christensen, U. R., Holzwarth, V., & Reiners, A. 2009, *Nature*, 457, 167
- Cowan, N. B. & Abbot, D. S. 2014, *ApJ*, 781, 27
- Cowan, N. B., Abbot, D. S., & Voigt, A. 2012a, *ApJ*, 752, L3
- Cowan, N. B. & Agol, E. 2011, *ApJ*, 729, 54
- Cowan, N. B., Agol, E., Meadows, V. S., et al. 2009, *ApJ*, 700, 915
- Cowan, N. B., Fuentes, P. A., & Haggard, H. M. 2013, *MNRAS*, 434, 2465
- Cowan, N. B., Machalek, P., Croll, B., et al. 2012b, *ApJ*, 747, 82
- Cowan, N. B., Robinson, T., Livengood, T. A., et al. 2011, *ApJ*, 731, 76
- Cowan, N. B. & Strait, T. E. 2013, *ApJ*, 765, L17
- Crossfield, I. J. M., Hansen, B. M. S., Harrington, J., et al. 2010, *ApJ*, 723, 1436
- Cushing, M. C., Roellig, T. L., Marley, M. S., et al. 2006, *ApJ*, 648, 614
- de Kok, R. J., Stam, D. M., & Karalidi, T. 2011, *ApJ*, 741, 59
- de Wit, J., Gillon, M., Demory, B.-O., & Seager, S. 2012, *A&A*, 548, A128
- Debes, J. H., Weinberger, A. J., & Schneider, G. 2008, *ApJ*, 673, L191
- Demory, B.-O., de Wit, J., Lewis, N., et al. 2013, *ApJ*, 776, L25
- Dent, W. R. F., Wyatt, M. C., Roberge, A., et al. 2014, *Science*, 343, 1490
- Dupuy, T. J. & Liu, M. C. 2012, *ApJS*, 201, 19
- Faherty, J. K., Rice, E. L., Cruz, K. L., Mamajek, E. E., & Núñez, A. 2013, *AJ*, 145, 2
- Faherty, J. K., Riedel, A. R., Cruz, K. L., et al. 2016, *ApJS*, 225, 10
- Faherty, J. K., Tinney, C. G., Skemer, A., & Monson, A. J. 2014, *ApJ*, 793, L16
- Farihi, J. 2016, *New A Rev.*, 71, 9
- Ford, E. B., Seager, S., & Turner, E. L. 2001, *Nature*, 412, 885

- Fortney, J. J., Lodders, K., Marley, M. S., & Freedman, R. S. 2008, *ApJ*, 678, 1419
- Fraine, J., Deming, D., Benneke, B., et al. 2014, *Nature*, 513, 526
- Fujii, Y. & Kawahara, H. 2012, *ApJ*, 755, 101
- Fujii, Y., Kawahara, H., Suto, Y., et al. 2011, *ApJ*, 738, 184
- Fujii, Y., Kawahara, H., Suto, Y., et al. 2010, *ApJ*, 715, 866
- Fujii, Y., Kimura, J., Dohm, J., & Ohtake, M. 2014, *Astrobiology*, 14, 753
- Gibson, N. P., Aigrain, S., Barstow, J. K., et al. 2013, *MNRAS*, 436, 2974
- Helling, C., Ackerman, A., Allard, F., et al. 2008, *MNRAS*, 391, 1854
- Hinkel, N. R., Timmes, F. X., Young, P. A., Pagano, M. D., & Turnbull, M. C. 2014, *AJ*, 148, 54
- Kaltenegger, L. & Sasselov, D. 2010, *ApJ*, 708, 1162
- Kao, M. M., Hallinan, G., Pineda, J. S., et al. 2016, *ApJ*, 818, 24
- Karalidi, T., Apai, D., Schneider, G., Hanson, J. R., & Pasachoff, J. M. 2015, *ApJ*, 814, 65
- Kasting, J. F., Whitmire, D. P., & Reynolds, R. T. 1993, *Icarus*, 101, 108
- Kataria, T., Showman, A. P., Fortney, J. J., Marley, M. S., & Freedman, R. S. 2014, *ApJ*, 785, 92
- Kawahara, H. 2016, *ApJ*, 822, 112
- Kawahara, H. & Fujii, Y. 2010, *ApJ*, 720, 1333
- Kawahara, H. & Fujii, Y. 2011, *ApJ*, 739, L62
- Kirkpatrick, J. D., Gelino, C. R., Cushing, M. C., et al. 2012, *ApJ*, 753, 156
- Knutson, H. A., Charbonneau, D., Allen, L. E., et al. 2007, *Nature*, 447, 183
- Knutson, H. A., Charbonneau, D., Cowan, N. B., et al. 2009, *ApJ*, 690, 822
- Knutson, H. A., Lewis, N., Fortney, J. J., et al. 2012, *ApJ*, 754, 22
- Kopparapu, R. k., Wolf, E. T., Haqq-Misra, J., et al. 2016, *ApJ*, 819, 84
- Kóspál, Á., Moór, A., Juhász, A., et al. 2013, *ApJ*, 776, 77

- Kreidberg, L., Bean, J. L., Désert, J.-M., et al. 2014, *Nature*, 505, 69
- Larsen, J. F. & Gardner, J. E. 2004, *Journal of Volcanology and Geothermal Research*, 134, 109
- Leconte, J., Forget, F., Charnay, B., Wordsworth, R., & Pottier, A. 2013, *Nature*, 504, 268
- Leggett, S. K., Cushing, M. C., Hardegree-Ullman, K. K., et al. 2016, ArXiv e-prints
- Liu, M. C., Magnier, E. A., Deacon, N. R., et al. 2013, *ApJ*, 777, L20
- Lodders, K. & Fegley, B. 2002, *Icarus*, 155, 393
- Macintosh, B., Graham, J. R., Barman, T., et al. 2015, *Science*, 350, 64
- Madhusudhan, N., Crouzet, N., McCullough, P. R., Deming, D., & Hedges, C. 2014, *ApJ*, 791, L9
- Majeau, C., Agol, E., & Cowan, N. B. 2012, *ApJ*, 747, L20
- Malhotra, R. 1993, *Nature*, 365, 819
- Marley, M. S. & Robinson, T. D. 2015, *ARA&A*, 53, 279
- Marois, C., Macintosh, B., Barman, T., et al. 2008, *Science*, 322, 1348
- Mason, B. G., Pyle, D. M., & Oppenheimer, C. 2004, *Bulletin of Volcanology*, 66, 735
- Meng, H. Y. A., Su, K. Y. L., Rieke, G. H., et al. 2014, *Science*, 345, 1032
- Metchev, S. A., Heinze, A., Apai, D., et al. 2015, *ApJ*, 799, 154
- Moór, A., Juhász, A., Kóspál, Á., et al. 2013, *ApJ*, 777, L25
- Morley, C. V., Fortney, J. J., Marley, M. S., et al. 2015, *ApJ*, 815, 110
- Morley, C. V., Marley, M. S., Fortney, J. J., et al. 2014, *ApJ*, 787, 78
- Mouillet, D., Larwood, J. D., Papaloizou, J. C. B., & Lagrange, A. M. 1997, *MNRAS*, 292, 896
- Oakley, P. H. H. & Cash, W. 2009, *ApJ*, 700, 1428
- O’Neill, C. & Lenardic, A. 2007, *Geophys. Res. Lett.*, 34, L19204

- O’Neill, C., Lenardic, A., Höink, T., & Coltice, N. 2013, *Mantle Convection and Outgassing on Terrestrial Planets*, ed. S. J. Mackwell, A. A. Simon-Miller, J. W. Harder, & M. A. Bullock, 473–486
- O’Neill, C., Lenardic, A., Weller, M., et al. 2016, *Physics of the Earth and Planetary Interiors*, 255, 80
- Pallé, E., Ford, E. B., Seager, S., Montañés-Rodríguez, P., & Vazquez, M. 2008, *ApJ*, 676, 1319
- Papale, P. & Polacci, M. 1999, *Bulletin of Volcanology*, 60, 583
- Parmentier, V., Showman, A. P., & Lian, Y. 2013, *A&A*, 558, A91
- Perrin, M. D., Duchene, G., Millar-Blanchaer, M., et al. 2015, *ApJ*, 799, 182
- Radigan, J., Jayawardhana, R., Lafrenière, D., et al. 2012, *ApJ*, 750, 105
- Radigan, J., Lafrenière, D., Jayawardhana, R., & Artigau, E. 2014, *ApJ*, 793, 75
- Rauscher, E. & Kempton, E. M. R. 2014, *ApJ*, 790, 79
- Rauscher, E. & Menou, K. 2013, *ApJ*, 764, 103
- Reiners, A. & Basri, G. 2008, *ApJ*, 684, 1390
- Robinson, T. D., Meadows, V. S., & Crisp, D. 2010, *ApJ*, 721, L67
- Robock, A., Adams, T., Moore, M., Oman, L., & Stenchikov, G. 2007, *Geophys. Res. Lett.*, 34, L23710
- Rodigas, T. J., Debes, J. H., Hinz, P. M., et al. 2014, *ApJ*, 783, 21
- Rogers, L. A. & Seager, S. 2010, *ApJ*, 712, 974
- Salyk, C., Pontoppidan, K. M., Blake, G. A., et al. 2008, *ApJ*, 676, L49
- Saumon, D. & Marley, M. S. 2008, *ApJ*, 689, 1327
- Schlichting, H. E. & Sari, R. 2007, *ApJ*, 658, 593
- Schlichting, H. E., Sari, R., & Yalinewich, A. 2015, *Icarus*, 247, 81
- Schneider, G., Grady, C. A., Hines, D. C., et al. 2014, *AJ*, 148, 59
- Scholz, A., Kostov, V., Jayawardhana, R., & Mužić, K. 2015, *ApJ*, 809, L29

- Schwartz, J. C. & Cowan, N. B. 2015, *MNRAS*, 449, 4192
- Schwartz, J. C., Sekowski, C., Haggard, H. M., Pallé, E., & Cowan, N. B. 2016, *MNRAS*, 457, 926
- Showman, A. P., Fortney, J. J., Lian, Y., et al. 2009, *ApJ*, 699, 564
- Showman, A. P. & Guillot, T. 2002, *A&A*, 385, 166
- Showman, A. P., Lewis, N. K., & Fortney, J. J. 2015, *ApJ*, 801, 95
- Simon, A. A., Rowe, J. F., Gaulme, P., et al. 2016, *ApJ*, 817, 162
- Sing, D. K., Fortney, J. J., Nikolov, N., et al. 2016, *Nature*, 529, 59
- Sing, D. K., Wakeford, H. R., Showman, A. P., et al. 2015, *MNRAS*, 446, 2428
- Skemer, A. J., Hinz, P. M., Esposito, S., et al. 2012, *ApJ*, 753, 14
- Skemer, A. J., Morley, C. V., Allers, K. N., et al. 2016, *ApJ*, 826, L17
- Snellen, I. A. G., Brandl, B. R., de Kok, R. J., et al. 2014, *Nature*, 509, 63
- Stark, C. C. & Kuchner, M. J. 2008, *ApJ*, 686, 637
- Sterzik, M. F., Bagnulo, S., & Palle, E. 2012, *Nature*, 483, 64
- Telesco, C. M., Fisher, R. S., Wyatt, M. C., et al. 2005, *Nature*, 433, 133
- Tingley, M. P., Stine, A. R., & Huybers, P. 2014, *Geophys. Res. Lett.*, 41, 7838
- Tremaine, S. 1991, *Icarus*, 89, 85
- Tsiganis, K., Gomes, R., Morbidelli, A., & Levison, H. F. 2005, *Nature*, 435, 459
- Valencia, D., O’Connell, R. J., & Sasselov, D. D. 2007, *ApJ*, 670, L45
- van Heck, H. J. & Tackley, P. J. 2011, *Earth and Planetary Science Letters*, 310, 252
- Wagner, K., Apai, D., Kasper, M., et al. 2016, *Science*, 353, 673
- Williams, D. M. & Gaidos, E. 2008, *Icarus*, 195, 927
- Wilson, L. 1980, *Journal of Volcanology and Geothermal Research*, 8, 297
- Wolf, E. T. & Toon, O. B. 2014, *Astrobiology*, 14, 241

- Wordsworth, R. D., Forget, F., Selsis, F., et al. 2011, *ApJ*, 733, L48
- Wyatt, M. C. 2003, *ApJ*, 598, 1321
- Wyatt, M. C. 2008, *ARA&A*, 46, 339
- Wyatt, M. C., Dermott, S. F., Telesco, C. M., et al. 1999, *ApJ*, 527, 918
- Yang, H., Apai, D., Marley, M. S., et al. 2015, *ApJ*, 798, L13
- Yang, J., Boué, G., Fabrycky, D. C., & Abbot, D. S. 2014, *ApJ*, 787, L2
- Yang, J., Cowan, N. B., & Abbot, D. S. 2013, *ApJ*, 771, L45
- Zhang, X. & Showman, A. P. 2014, *ApJ*, 788, L6
- Zhou, Y., Apai, D., Schneider, G. H., Marley, M. S., & Showman, A. P. 2016, *ApJ*, 818, 176
- Zugger, M. E., Kasting, J. F., Williams, D. M., Kane, T. J., & Philbrick, C. R. 2010, *ApJ*, 723, 1168
- Zugger, M. E., Kasting, J. F., Williams, D. M., Kane, T. J., & Philbrick, C. R. 2011, *ApJ*, 739, 12