


SAG15 DRAFT REPORT

SCIENCE QUESTIONS FOR DIRECT IMAGING EXOPLANET MISSIONS



DANIEL APAI AND THE SAG15 TEAM

October 6, 2016

Contents

| | | |
|----------|---|-----------|
| 1 | The SAG15 Team and Contributors | 3 |
| 2 | Introduction | 4 |
| 3 | Overview of Science Questions | 4 |
| 4 | Exoplanetary System Characterization | 6 |
| 4.1 | A1. What is the diversity of planetary architectures? Are there typical classes/types of planetary architectures? How common are Solar System-like planetary architectures? | 6 |
| 4.1.1 | Complementary Non-Imaging Data | 8 |
| 4.2 | A2. What are the distributions and properties of planetesimal belts and exozodiacal disks in exoplanetary systems and what can these tell about the formation and dynamical evolution of the planetary systems? | 9 |
| 4.2.1 | Current Knowledge | 9 |
| 4.2.2 | Sub-questions | 10 |
| 4.2.3 | Complementary Data | 11 |
| 5 | Exoplanet Characterization | 13 |
| 5.1 | B1. How do rotational periods and obliquity vary with orbital elements and planet mass/type? | 13 |
| 5.1.1 | State of the Art to Measure Rotational Periods | 13 |
| 5.1.2 | Science Cases | 15 |
| 5.1.3 | The Science Value of Independently Measured Planet Masses and Radii | 18 |
| 5.2 | B2: Which rocky planets have liquid water on their surfaces? Which planets have continents and oceans? | 26 |
| 5.2.1 | Detecting Oceans | 26 |
| 5.2.2 | Liquid Water Clouds | 28 |

| | | |
|----------|--|-----------|
| 5.3 | B3. What are the origins and composition of condensate clouds and hazes in ice/gas giants and how do these vary with system parameters? | 30 |
| 5.3.1 | Science Value of Independently Measured Planet Masses and Radii | 36 |
| 5.4 | B4. How do photochemistry, transport chemistry, surface chemistry, and mantle outgassing effect the composition and chemical processes in terrestrial planet atmospheres (both habitable and non-habitable)? | 37 |
| 5.4.1 | Science Value of Independently Measured Planet Masses and Radii | 38 |
| 6 | Exoplanetary Processes | 39 |
| 6.1 | C1. What processes/properties set the modes of atmospheric circulation and heat transport in exoplanets and how do these vary with system parameters? | 39 |
| 6.2 | C2. What are the key evolutionary pathways for rocky planets and what first-order processes dominate these? | 45 |
| 6.2.1 | Science Value of Independently Measured Planet Masses and Radii | 46 |
| 6.3 | C3. What types/which planets have active geological activity, interior processes, and/or continent-forming/resurfacing processes? | 47 |
| 6.3.1 | Geological Activity and Plate Tectonics on Extrasolar Rocky Planets | 49 |
| 6.3.2 | Observational Methods | 50 |
| 6.3.3 | Complementary Datasets | 52 |
| 6.3.4 | Science Value of Independently Measured Planet Masses and Radii: Very High | 52 |
| 7 | Data Requirements | 53 |
| A | SAG15 Charter | 54 |

1. The SAG15 Team and Contributors

Chair: Daniel Apai, University of Arizona (apai@arizona.edu)

Members:

| | |
|--|---|
| Travis Barman, University of Arizona | Patrick Lowrence, IPAC/Caltech |
| Alan Boss, Carnegie DTM | Avi Mandell, NASA GSFC |
| James Breckenridge, Caltech | Mark Marley, NASA Ames |
| David Ciardi, IPAC/Caltech | Michael McElwain, NASA GSFC |
| Ian Crossfield, University of Arizona | Nikku Madhusudhan, Cambridge University |
| Nicolas Cowan, McGill University | Charley Noecker, JPL |
| William Danchi, NASA GSFC | Ilaria Pascucci, University of Arizona |
| Shawn Domagal-Goldman, NASA GFSC | Peter Plavchan, Missouri State University |
| Yuka Fujii, NASA GISS | Aki Roberge, NASA GSFC |
| Caroline Morley, Lick Observatory | Leslie Rogers, University of Chicago |
| Glenn Schneider, University of Arizona | Adam Showman, University of Arizona |
| Nicolas Iro, University of Hamburg | Arif Solmaz |
| Stephen Kane, San Francisco State University | Philip Stahl, NASA MSFC |
| Theodora Karalidi, University of Arizona | Karl Stapelfeldt, JPL |
| Markus Kasper, ESO | Mark Swain, JPL |
| James Kasting, Penn State University | Margaret Turnbull, SETI Institute |
| Ravikumar Kopparapu, NASA GSFC | William Moore |
| | Anthony del Genio, NASA GISS |
| | Eric Mamajek, JPL |

2. Introduction

This report presents organized input from the international exoplanet community on science questions that can be answered by direct imaging missions.

For each science question we also explore the types and quality of datasets that are either required to answer the question or greatly enhance the quality of the answer. We also highlight questions that require or benefit from complementary (non-direct imaging) observations.

In preparing the report no specific mission architecture or requirements were assumed or advocated for; however, where obvious connections to planned or possible future mission existed these were identified. More detailed evaluations of the capabilities of specific mission architectures are provided in other SAG reports and by ongoing NASA STDs studies. The SAG15 report does not include discussion of biosignatures or planets transformed by life, which are discussed in the ongoing SAG16 study, however, the SAG15 reports does include discussion of the characterization of habitable zone earth-sized planets.

Community input: Input for this report has been collected and comments on the different report drafts have been solicited through a range of channels, including: i) SAG15 website (<http://eos-nexus.org/sag15>); ii) monthly SAG15 telecons; iii) breakout and discussion sessions during related workshops and meetings; iv) direct requests from topical experts; v) email invitations and solicitations via the EXOPAG and NExSS mailing lists.

Author list and contributor list: The final report will represent the full endorsement of each author, based on their explicit written statements. In contrast, the SAG15 team list and list of contributors provided in the interim drafts only represents experts who provided input or joined the SAG15 team. The contributor list in the report drafts, therefore, does not represent the endorsement of the draft report and its findings by the contributors.

3. Overview of Science Questions

The science questions in this report are divided into three categories (see Table 2). Questions in Category A aim at the statistical characterization of the formation, evolution, and properties of planetary systems. Questions in Category B aim at the quantitative characterization of individual planets or small groups of planets. Questions in Category C aim at understanding processes that shape planets and planetary atmospheres through comparative studies of planets.

Table 1: Overview of the science questions discusses in this report.

| High-Level Science Questions | Priority |
|--|----------|
| Science questions on exoplanetary system architectures and population | |
| A1. What is the diversity of planetary architectures? Are there typical classes/types of planetary architectures? How common are planetary architectures resembling the Solar System? | |
| A2. What are the distributions and properties of planetesimal belts and exo-zodiacal disks in exoplanetary systems and what can these tell about the formation and dynamical evolution of planetary systems? | |
| Science questions on exoplanet properties | |
| B1. How do rotational periods and obliquity vary with orbital elements and planet mass/type? | |
| B2. Which rocky planets have liquid water on their surfaces? | |
| B3. What are the origins and composition of clouds and hazes in ice/gas giants and how do these vary with system parameters? | |
| B4. How do photochemistry, transport chemistry, surface chemistry, and mantle outgassing affect the composition and chemical processes in terrestrial planet atmospheres (both habitable and non-habitable)? | |
| Science questions on exoplanet evolution and processes | |
| C1. What processes/properties set the modes of atmospheric circulation and heat transport in exoplanets and how do these vary with system parameters? | |
| C2. What are the key evolutionary pathways for rocky planets? | |
| C3. What types/which planets have active geological activity, interior processes, and/or continent-forming/resurfacing processes? | |

4. Exoplanetary System Characterization

4.1. A1. What is the diversity of planetary architectures? Are there typical classes/types of planetary architectures? How common are Solar System-like planetary architectures?

Contributors: Daniel Apai, Nicholas Cowan, Renyu Hu

Suggested referees: John Johnson, Tim Morton

The term *planetary system architecture* is used here as a descriptor of the high-level structure of a planetary system as given by the stellar mass, the orbits and mass of the planets, as well as the location and mass of its planetesimal belts.

Understanding the diversity of planetary architectures is important for at least the following two reasons: i) The diversity of planetary system architectures is expected to reflect *the range of possible formation and evolution pathways* of planetary systems. ii) To understand how common true *Earth analogs* are we must understand *how common are planetary systems with architectures similar to that of the Solar System*.

Our current picture of planetary system architectures builds on five sources: 1) Solar System; 2) Data from transiting exoplanets, primarily the Kepler Space Telescope, which probe the inner planetary systems (typically up to periods of 1 year); 3) radial velocity surveys, which provide data on planets with masses typically larger than those accessible to Kepler observations, but over multi-year periods; 4) microlensing surveys, which are also sensitive to small rocky planets at intermediate periods, but provide as yet limited statistics; 5) direct imaging surveys: capable of probing giant exoplanets at semi-major axes of 8 au or greater.

Based on the extrapolation of the close-in exoplanet population detected by the Kepler mission it is very likely that we do not yet have an efficient planet detection method to sample the *majority* of exoplanets that exists (at intermediate to large periods and with masses comparable to Earth). ESA’s Gaia mission will increase the census of known intermediate-to long-period giant planets by about $\sim 3,000$ new discoveries. In addition, the proper motion information for the Solar neighborhood will improve the identification and age-dating of co-moving stellar groups which, in turn, will greatly reduce the uncertainties in the giant planet mass-to-luminosity conversion used by ground-based direct exoplanet imaging surveys, thereby improving the long-period giant planet occurrence rate and mass distribution measurements.

Furthermore, the gradually extending baselines and improving accuracy of radial velocity measurements will also further improve the occurrence rates for short and intermediate-orbit planets (most significantly for neptune-mass and larger planets). In spite of these significant improvements the occurrence rates of the sub-neptune planets (including rocky and icy planets) at intermediate- to long-period orbits is presently poorly known and will remain largely unconstrained in the near future.

A direct imaging mission would be powerful in surveying low-mass planets at intermediate and long orbits (~ 1 to 30 au), establishing their orbits or constraining their orbital parameters, and measuring or deducing their masses and sizes.

Although different techniques will sample different planet populations around different set of stars, a capable direct imaging mission can have the capability of providing a more complete census of planets in the targeted systems than current methods. Direct imaging will survey planets in a range of orbital distances from their host stars, determined by the planet-star separation and photometry contrast. In addition, multiple visits are required to build a more complete census in the search range of orbital distances, because of the planets' changing orbital phase (Greco & Burrows 2015).

Sub-questions:

- *What is the diversity of planetary architectures?* The statistical assessment of the occurrence rate and mass distribution of planets as a function of system parameters (e.g., stellar mass, composition) can constrain and/or verify the predictions of planet formation models. The dispersion in different parameters (from data corrected for selection effects and biases) can be used to quantify the diversity of the architectures.
- *Are there typical classes/types of planetary architectures?* If there are different typical planet formation or evolution pathways, these may lead to the emergence of different classes of planetary architectures (e.g., planetary systems with hot jupiters). The presence of classes of planetary systems may be identified as clustering in the multi-dimensional parameter space that describes planetary architectures.
- *How common are Solar System-like planetary architectures?* The *local density* of the systems in the multi-dimensional parameter space that describes planetary architectures, determined at the location of the Solar System provides a measure of the occurrence rate of Solar System-like architectures. Furthermore, in this multi-dimensional parameter space distance-type metrics can be defined to reflect the similarity of any two planetary systems. Although non-unique, such metrics may be used to explore the frequency of systems as a function of distance from the Solar System to establish which nearby systems are the most similar to ours.

4.1.1. Complementary Non-Imaging Data

- *Radial velocity:* Constraints from radial velocity measurements can greatly reduce the number of direct imaging epochs required to establish the orbital elements of the planets. These measurements can also constrain or determine the mass of the target planets.
- *Microlensing:* Statistical constraints from the WFIRST-Microlensing survey will provide important context for the frequency of medium-separation low-mass planets.
- *Ground-based adaptive optics imaging:* These observations may will be capable of discovering super-earth, neptune, and gas giant exoplanets at intermediate and long periods in nearby systems. By providing positions at additional epochs they will place constraints on the orbits of the planets. In combination with Gaia astrometry the planetary architectures of nearby systems will be relatively well explored in 10-20 years.
- *Gaia Astrometry:* This dataset will provide orbital elements and masses for a large number of intermediate- to long-period gas giant planets, an important statistical context for the planets to be discovered by the direct imaging mission.

Observational Requirements:

Sample size: Medium to large samples (30 to 100).

Observations/Data: Optical or infrared imaging to identify the presence and location of planets in each system. Multi-epoch imaging (or complementary radial velocity or astrometry) is required to constrain orbital parameters.

Pending Questions / Comments:

- 1) How many epochs are required to establish orbital parameters?
- 2) To what accuracy should the orbital parameters be measured to?
- 3) What statistical constraints will WFIRST-ML, Gaia, and future RV surveys provide?
- 4) Add more info on GMT/ELT/TMT capabilities.

4.2. A2. What are the distributions and properties of planetesimal belts and exo-zodiacal disks in exoplanetary systems and what can these tell about the formation and dynamical evolution of the planetary systems?

Contributors: Daniel Apai

Direct imaging missions will provide spatially resolved images of exo-zodiacal disks, possibly composed of narrow and/or extended dust belts. In these belts dust is produced by minor body collisions and the dust belts are dynamically sculpted by the gravitational influence of the star and the planets, grain-grain collisions, as well as radiation pressure (for reviews see, e.g., Wyatt 2008). In some, apparently very rare, systems gas is also present and may influence the dust distribution.

The distribution and properties of exo-zodiacal dust belts (or debris disks) are important as they provide information on:

- The presence, orbits, and masses of unseen planets orbiting in the disks.
- The orbits and masses of planets seen in the direct images, but for which orbits are not known.
- The inclination of the disk/planet system.
- The formation and evolution of the system, including the past migration and orbital rearrangements of the planets.
- Compositional constraints on the availability of volatiles/organics in the planetesimal belts and, by inference, in the planets.

4.2.1. Current Knowledge

Currently, large databases of bright debris disks are available for which spatially unresolved thermal infrared observations (spectral energy distributions or SEDs) are available. In addition, for a subset of disks spatially resolved scattered light or thermal emission images

are available (see, e.g., Fig. 1). Mid-infrared spectroscopy of solid state dust features (e.g., Telesco et al. 2005) and polarimetric imaging provide additional constraints on dust composition and disk structure (e.g., Perrin et al. 2015). The different wavelengths, the types of emission (continuum, spectral features), and the polarization properties of the light allow us to disentangle the different dust components and study their origins.

Add a few sentences on general debris disk evolution, collisional events, self-stirring and planetary stirring, orbital rearrangements and bombardments.

4.2.2. Sub-questions

The presence, orbits, and masses of unseen planets: Detailed simulations of debris disk structures and disk-planet interactions provide predictions for the expected disk structures (see Fig. 2, e.g., Wyatt et al. 1999; Wyatt 2003; Mouillet et al. 1997; Stark & Kuchner 2008). In a large set of disks complex structures have been observed which can possibly be explained by the influence of yet unseen planets (e.g., Schneider et al. 2014); in a very small number of systems disks and planets have been observed together, providing an opportunity to study disk-planet interactions and to validate models (see, e.g., Apai et al. 2015 and Fig. 1).

The orbits and masses of planets seen in the direct images: With certain direct imaging architectures (e.g., starshades) opportunities for multi-epoch observations may be limited, making it more difficult to verify that point sources are planets and not background sources; and to estimate masses/orbits for the planets from short integrations. Most directly imaged systems are expected to host dust disks, whose structures may be used to verify that the planet candidates imaged are indeed in the system and then to constrain their mass and orbit.

The inclination of the disk/planet system: For any planet an important but particularly challenging parameter pair to determine is the inclination/eccentricity pair. These quantities are partially degenerate and can be difficult to disentangle from observations limited to a handful of visits. Resolved debris disks structures can complement measurements of the planet’s relative motion to break the degeneracy of inclination/eccentricity. For example, nearly-edge on disks can be recognized even in single-epoch images, which then greatly constrain the available parameter space for the planet’s orbit.

The formation and dynamical evolution of the systems: The mass and position of planetesimal belts can provide powerful constraints on the formation and evolution of planetary systems, including planet migration and/or major orbital rearrangements. For example, the asteroid belt and the Kuiper belt in the Solar System have revealed such orbital rearrange-

ment and potential past instabilities (e.g., Malhotra 1993; Tsiganis et al. 2005). In addition, sensitive time-resolved observations in debris disks also have the potential to identify multiple other mechanisms that act on short timescales, such as the aftermath of recent major impacts (e.g., Meng et al. 2014), dust clumps moving under the influence of radiation pressure, or dust created by planetesimals trapped in resonant structures (e.g., Wyatt 2003; Apai et al. 2015; Boccaletti et al. 2015).

Compositional constraints on the availability of volatiles/organics in the planetesimal belts: In each system planetesimal belts are leftover reservoirs of the same material that formed the planets and therefore the planetesimal’s composition constrains the composition of the planets themselves. Of particular interest is the availability of volatiles and organics in the planetesimals, as these are thought to be heavily depleted in the warm, inner disk regions where habitable planets accrete. Organics and volatile content (interior or as a surface layer) change the optical properties of the dust grains, producing signatures that are detectable at optical and infrared wavelengths (e.g., Debes et al. 2008; Rodigas et al. 2014; Ballering et al. 2016). Recently discovered debris disks with gas content that may be recent or primordial provide an additional opportunity to explore volatile reservoirs in planetesimal belts (Dent et al. 2014; Kóspál et al. 2013; Moór et al. 2013)

4.2.3. Complementary Data

Exo-zodiacal disk studies will benefit from:

- 1) WFIRST imaging of debris disks;
- 2) ALMA observations of cold debris disks;
- 3) LBTI observations of the warm debris;
- 4) JWST observations of warm debris disks.

Observational Requirements

Sample size: Individual or small samples are useful for the characterization of individual disk-planets systems, but medium to larger samples required for studying disk evolution or the distributions of disk properties.

Observations: Multi-wavelength optical/near-infrared/mid-infrared imaging, spectroscopy, and optical/near-infrared polarimetry. Large field of view ($\sim 10''$ or larger) may be needed to study disk morphology in nearby systems.

Questions to SAG15:

Input from WFIRST PS team on what debris disk science do they foresee.

5. Exoplanet Characterization

5.1. B1. How do rotational periods and obliquity vary with orbital elements and planet mass/type?

Contributors: Daniel Apai, Nicolas Cowan, Renyu Hu, Anthony del Genio **Suggested referees:** Hajime Kawahara

A planet’s rotational state refers to both its obliquity and frequency, or equivalently period. Planetary rotation constrains the formation and angular momentum evolution of a planet, especially when comparing statistical samples of diverse planets. Moreover, the rotation of a given planet impacts its climate through diurnal forcing and through the Coriolis forces, and contributes to magnetic field generation.

For example, Yang et al. (2014, 2013) showed that the rotation periods of temperate terrestrial planets changes the inner boundary of the habitable zone by a factor of two in insolation (also see Kopparapu et al. 2016). Furthermore, planetary magnetic fields may be important shields against atmospheric loss. As these examples illustrate the rotational state of temperate terrestrial planets directly impacts their habitability.

We note, that depending on the nature and atmospheric composition of a planet its true rotational period (that of its bulk mass) may or may not be possible to determine observationally. For example, while a rocky planet’s rotational period may be observed via the observations of surface features, for gaseous planets or rocky planets with optically thick atmospheres the rotational period of the interior may remain hidden and only an ”apparent rotational period” may be observed: one that is a combination of the rotational rate and dominant atmospheric motions (winds, circulation).

5.1.1. *State of the Art to Measure Rotational Periods*

As of now little is known about the obliquity and rotational periods of non-synchronously rotating exoplanets. Rotational periods for planets and exoplanets have been determined through four different methods:

a) Phase Curve for Irradiated Planets: For some close-in synchronously rotating giant exoplanets the orbital/rotational phase modulation is detectable in the combined light of the star and planet system. The modulation allows coarse two-dimensional mapping of the planets: For example, the dayside map of HD 189733b suggests that this hot Jupiter has zero obliquity (Majeau et al. 2012; de Wit et al. 2012). The eastward offset of the hotspot

observed on most hot Jupiters (Knutson et al. 2007, 2009, 2012; Crossfield et al. 2010; Cowan et al. 2012b) is consistent with equatorial super-rotation on a synchronously-rotating planet (Showman & Guillot 2002), but also with slower winds on a non-synchronous planet (Rauscher & Kempton 2014). In fact, there is a complete observational degeneracy between the rotation of a gaseous exoplanet and its winds (Cowan & Agol 2011). (Yang et al. 2013) showed that tidally locked temperate planets will have dayside ??? at 220 W/m^2 , while non-synchronously rotating but otherwise similar planets will not.

b) Period of the magnetic field’s rotation: The magnetic field is tracing the rotational periods of the planets’ *interiors*, which may be different from the latitude-averaged rotational periods measured in their upper atmospheres. In the Solar System, Jupiter’s rotational period is defined by the rotation of its inclined (w.r.t. spin axis) magnetic dipole, while Saturn’s magnetic field exhibits a very small tilt and its rotation period thus remains somewhat uncertain. Recent detections of modulated radio emission from nearby brown dwarfs (e.g., Kao et al. 2016) suggest that very sensitive radio-wavelength observations of extrasolar giant planets may also be used in the future to establish their rotational periods.

c) Absorption line width measurements: Recently, CO absorption line width measurements have been used to measure the rotational velocity ($v \sin i$) for the directly imaged exoplanet Beta Pictoris b (Snellen et al. 2014) and in the combined star and planet light for hot Jupiters (e.g., (Rodler et al. 2012)). Similar studies for rotational line broadening have been carried out successfully for brown dwarfs (e.g., Reiners & Basri 2008). In order to convert the observed $v \sin i$ into a rotational period, one must know the planet’s radius and obliquity. This method is therefore well-suited for brown dwarfs and giant planets (which are all approximately the size of Jupiter), but could prove problematic for lower-mass directly-imaged planets of unknown radius. Furthermore, it is more applicable for systems where constraints exist on the planets’ obliquities (primarily derived from rotational modulations observed over multiple orbital phase angles).

d) Rotational photometric/spectroscopic modulations in hemisphere-integrated light for directly imaged exoplanets (Fig. 4, Zhou et al. 2016) and planetary-mass brown dwarfs (e.g., Biller et al. 2015, Leggett et al. 2016). This method is conceptually identical to method *a*, but requires a different observational approach. Brown dwarfs (planetary mass and more massive), are good analogs for directly imaged exoplanets. These observations showed that low-level ($\sim 1\%$) rotational modulations in thermal emission are *very* common (Buenzli et al. 2014; Metchev et al. 2015), and can be used to measure or constrain rotational periods and to study cloud properties (e.g., Artigau et al. 2009; Radigan et al. 2012; Apai et al. 2013). Similarly, reflected-light observations of Solar System giant planets have also been used to demonstrate that rotational periods and their cloud covers can be characterized

(e.g., Jupiter: Karalidi et al. 2015; Neptune: Simon et al. 2016).

Both techniques *a* and *d* may be applicable for exoplanets directly imaged with next-generation space telescopes. While method *b* requires high spectral resolution and provides Doppler information, method *c* requires only high signal-to-noise time-resolved photometry and not strongly wavelength-dependent.

5.1.2. *Science Cases*

Habitable Planets (Earth-sized and Super-Earths): Rotation rates are an important parameter for climate and atmospheric circulation models of habitable planets: they constrain diurnal temperature modulations, determine the strength of the Coriolis force, the nature of the circulation, and thus the location of clouds, influence current and past magnetic field strengths and geometry, and indirectly constrain the atmospheric loss that may have occurred on these planets. Comparative studies of dynamo-generated magnetic energy densities in Solar System planets, the Sun, and rapidly-rotating low-mass stars show a correlation between the magnetic field strengths and the density and bolometric flux of the objects (see Fig. 5, e.g., Christensen et al. 2009). These studies argue for a scaling relation, based on Ohmic dissipation, where the field strength is only weakly sensitive to rotation rate, but the rotational rate fundamentally impacts the magnetic field geometry (bipolar vs. multi-polar, Christensen 2010). Furthermore, rotational rates also carry information about the accretion history of the planets and, in particular, about the size distribution of the planetary building blocks (e.g., Schlichting & Sari 2007).

In addition, the obliquity of habitable planets also has a major impact on the seasonal and diurnal temperature variations and on their climate in general. Obliquity is much more difficult to determine than the rotational rate. However, simulated observations demonstrate that it is possible to determine this quantity from high signal-to-noise reflected light lightcurves obtained at multiple orbital phases.

Considerable effort was put into exploring time-resolved observations of Earth, as exoplanet analog. Researchers have used simulated disk-integrated brightness variations of Earth to demonstrate that its rotational period can be estimated, even in the presence of time-varying clouds (Pallé et al. 2008; Oakley & Cash 2009). Likewise, such observations spanning multiple orbital phases constrain obliquity (Kawahara & Fujii 2010, 2011; Fujii & Kawahara 2012; Schwartz et al. 2016; Kawahara 2016). Schwartz et al. (2016) showed that although both latitudinal and longitudinal heterogeneities contribute to the obliquity signal, the latter contains more information. In principle, the amplitude modulation of rotational

variations at only three orbital phases uniquely identifies a planet’s obliquity vector (the obliquity and its orientation with respect to the observer’s line of sight). Taking the complementary *frequency modulation* approach, Kawahara (2016) showed that modest signal-to-noise observations spanning most of a planet’s orbit could also constrain a planet’s obliquity, even if one is agnostic of the planet’s albedo map. A comprehensive study by Schwartz et al. (2016) demonstrated that planetary obliquity can be constrained from observations at just a few orbital phase angles (see Figure 6).

A special case of rocky planets are those with very thin or no atmosphere (analogous to a “super-Mars” or a “dry Earth”, an Earth-like planet that formed essentially dry or lost its atmosphere and water). Such planets may form as a result of extensive atmospheric loss due to evaporation (Hot super-Mars), stellar wind stripping, or impact stripping (e.g., Schlichting et al. 2015). At pressures lower than water’s triple point (6 mbar) liquid water is not stable, even if the planet is otherwise Earth-sized and it is inside the habitable zone. The ability to measure rotational periods for these planets may provide important insights into the mechanism that led to the complete atmospheric loss. Atmosphereless are suitable for direct measurements of their rotational periods, because various types of rocky surfaces (i.e., mineral assemblages) have deep and wide albedo features that will introduce photometric rotational modulations in the visible and near-infrared (Hu et al. 2012a).

Fujii et al. (2014) used albedo-map generated lightcurves and, where available, observed photometric variations to explore the geological features detectable on diverse Solar System bodies with minor or no atmospheres (Moon, Mercury, the Galilean moons, and Mars). The study included the evaluation of the light curves and the features that are detectable at wavelengths ranging from UV through visible to near-infrared wavelengths, and also explored the accuracy required to determine the rotational periods of these bodies. Figure 7 provides an example for the wavelength-dependence of the rotational variability amplitudes in different bodies.

Gas and Ice Giant Exoplanets: The rotational periods of gas/ice giants may also be useful for constraining their formation and evolution (Tremaine 1991) and important for understanding their atmospheric circulation. Non-axisymmetrically distributed condensate clouds and hazes (photochemical or other origin) will introduce rotational modulations, both in reflected and in thermal emission (e.g., Simon et al. 2016). In addition, polarimetric modulations introduced by light scattering on heterogeneously distributed dust/haze grains may also be detectable. Currently, rotational rate estimates exist for close-in exoplanets (assumed to be equal to their orbital periods) and a few measurements exist for directly imaged exoplanets and planetary-mass brown dwarfs. The rotational angular momenta of close-in exoplanets (i.e., synchronously rotating) is reset by tidal interactions and no longer

carries information on the intrinsic angular momenta of the objects. In contrast, the angular momenta of non-synchronously rotating exoplanets (such as those probed via direct imaging) carry information about their formation and angular momentum evolution. Photometric modulations have been measured in two near-infrared filters for the $\sim 4\text{--}6\text{ M}_{Jup}$ exoplanet 2M1207b (Zhou et al. 2016) and led to a rotational period measurement of $10.7_{-0.6}^{+1.2}$ h. CO absorption line rotational broadening measurements for the $10\text{--}13\text{ M}_{Jup}$ planet β Pictoris b suggests a $v \sin i = 15$ km/s, which – assuming an equatorial viewing geometry, age, and mass – suggests a very similar rotational period. Similarly to these young exoplanets, photometric variations were used to measure the rotational periods of unbound young planetary mass-objects (Biller et al. 2015; Leggett et al. 2016) and very low-mass brown dwarfs (e.g., Scholz et al. 2015). The picture emerging – based on the very limited data – suggests that super-jupiter exoplanets and low-mass brown dwarfs start with similar angular momenta and during their evolution (cooling and contraction) their rotation rate increases, converging to the extrapolation of the Solar System mass-period relationship (see Figure 4).

A direct imaging mission capable of obtaining moderately high signal-to-noise ratio photometry of giant exoplanets can study possible trends between planet mass, semi-major axis, and rotational period.

Obliquity for gas giants: For gas giants (with well-constrained radii) combining the rotational period determined from rotational modulations with radial velocity information (line broadening due to rotation) allows constraining or deriving the rotation and inclination of the planet (e.g., Allers et al. 2016). Finally, the Fourier spectrum or polarimetry of thermal emission (de Kok et al. 2011; Cowan et al. 2013) as well as the amplitude and frequency modulation of reflected light rotational variations can provide an obliquity estimate (Schwartz et al. 2016; Kawahara 2016).

A Note on Hazy Atmospheres: Planets with thick haze layers may pose a challenge for rotational signals using methods c and d (line width measurements and temporal photometric/spectroscopic variations) depending on the wavelengths of observations and the origins of molecular absorption or cloud features studied). Because haze particles *by definition* are small ($\sim 0.01\text{--}1\text{ }\mu\text{m}$) and are not modulated by large-scale condensation-evaporation patterns associated with vertical motions the way clouds are, they sediment more slowly and their residence time in the atmosphere will be much longer than the rotational period ($t_{res} \gg P$). This may result in featureless haze layers (e.g., Venus), unless other absorbing constituents that are sensitive to the atmospheric circulation are present. As haze particles can be generated at higher altitudes than larger particles produced by condensation, the featureless haze layers *if optically thick* will mask any heterogenous condensate cloud structure as well as any surface structures. Similarly, optically thick haze layers may cover or weaken

the rotationally broadened line profiles in the atmospheres, also limiting the use of Doppler techniques. Therefore, planets enshrouded in thick haze layers may often not be well suited for rotational studies.

The two hazy planets in our solar system are useful cases in point. Venus is shrouded in a $\sim 1\mu\text{m}$ sulfuric acid haze but with dark ultraviolet features due to an unknown absorber that revealed a ~ 4 -day rotational period in ground-based observations (Boyer & Camichel 1961; Traub & Carleton 1975). This was later shown to be due to the atmosphere’s superrotation rather than the slow 243 day rotation period of its surface (Rossow et al. 1990). Titan is covered by a stratospheric hydrocarbon haze that is featureless except for a seasonally varying hemispheric albedo asymmetry (e.g., Lorenz et al. 2009). The haze obscures the view of tropospheric methane clouds and the surface, but these can be detected in near-infrared imagery (Turtle et al. 2011; Rodriguez et al. 2011).

5.1.3. *The Science Value of Independently Measured Planet Masses and Radii*

Planetary Radius: For methods that measure rotational velocity rather than period, knowledge of planetary radius and obliquity is required to convert rotational broadening into a rotational period. However, if the goal is to determine the Coriolis forces, then rotational broadening is sufficient. For the photometric methods that produce a period estimate, on the other hand, the frequency of diurnal forcing is easily derived, while estimating the Coriolis forces again requires the planetary period. In general, rotational information is most useful when combined with radius estimates. No complementary observations are required for science results from rotational period measurements, but observations constraining the planetary orbits may be combined with the obliquity and rotational period to constrain the formation history of low-mass planets.

Giant Planets: Radii for mature giant planets will be close to one Jupiter radius, but masses may vary by an order of magnitude. Masses may be derived from spectral retrieval that includes a fit for surface gravity.

Radii and masses of rocky planets vary *more* than those of giant planets: mass may vary by a factor of ~ 20 (from Mars to super-Earths): while rotational periods alone will be important and useful for atmospheric circulation models, mass and/or radius measurements would yield important additional science: mass measurements would allow exploring trends between formation mechanisms and angular momentum; and radius estimates (even from mass-radius relationships) would allow calculating Coriolis forces from rotational periods, significantly constraining the atmospheric circulation models.

Planet mass measurements from radial velocity or astrometry, or gravitational interactions between the planets, can be combined with rotational periods to determine the angular momenta of the giant planets, which may be useful for constraining their accretion history.

The periodicity in photometric variations is a direct measure of the rotational period, i.e., rotational period measurements do not require mass measurements. However, verifying the predicted trend between angular momentum, orbital period, mass (which potentially constrains the formation history) requires mass and radius measurements.

Observational Requirements (draft)

Sample size: Individual to medium/large

Observations: Very high spectral resolution; OR multi-epoch photometry.

Obliquity: photometry over complete rotational phase at multiple distinct orbital phases (at minimum three phases).

Pending Comments/Questions:

Kasper: ELT HCI + HRS will allow us to study rotation periods, sample will be small
- not sure about statistical relevance

Table 2: Expected rotational modulation amplitudes and constraints on rotational period and obliquity for terrestrial and giant exoplanets.

| Planet Type | Optimal λ | Amplitude | Acceptable λ | Amplitude | Baseline |
|--------------------------|-------------------|-----------|-----------------------|-----------|--------------|
| <i>Rotational Period</i> | | | | | |
| Terrestrial | 0.9 μm | 25% | 0.5-10 μm | 10–35% | P=3–30 h |
| Ice/Gas Giant | 5 μm | 15% | 0.3-5.0 μm | 3% | P=3–20 h |
| <i>Obliquity</i> | | | | | |
| Terrestrial | 0.9 μm | 25% | 0.5-10 | 10–35% | 3 \times P |
| Ice/Gas Giant | 5 μm | 15% | 0.3-5.0 μm | 3% | 3 \times P |




Figure 14. Key structures in the β Pic system, as derived from multi-wavelength imaging.




Fig. 1.— Simulations of the structure of the edge-on debris disk around Beta Pictoris correctly predicted the location and mass of the perturber super-Jupiter Beta Pictoris b (Mouillet et al. 1997). This system is one of the the best-studied examples of disk-planet interactions. *Lower panel:* HST/STIS coronagraphic image (blue), ALMA dust continuum (green), and ALMA CO gas emission (red) illustrate the complex structure of the disk (from Apai et al. 2015).




Fig. 2.— Comparison of the optical depths predicted by disk-planet interactions models for a composite cloud formed for a 2 earth-mass planet at 6 au (from Stark & Kuchner 2008). The planet, marked with a white dot, orbits counterclockwise in these images. *Left*: Optical depth of the smallest particles included in the composite clouds. *Right*: Optical depth of the largest particles included in the composite clouds. The largest particles dominate the optical depth in a cloud of particles released with a Dohnanyi crushing law, because they are longer-lived and more likely to be trapped in mean motion resonances than smaller particles.




Fig. 3.— Whitened power spectrum from 50-day-long Kepler monitoring of hemisphere-integrated reflected light Neptune, with the most significant peak corresponding to the rotation period. Numbers above some peaks indicate the latitudes on Neptune corresponding to that rotation period based on the zonal velocities. From Simon et al. (2016).




Fig. 4.— Rotation periods provide insights into the properties and formation of planets. A comparison of Solar System planets, directly imaged exoplanets, and brown dwarfs reveals a characteristic mass-dependent rotation rate for massive planets. The ages of the Solar System planets is 4.56 Gyr; the ages of the directly imaged planets is <30 Myr; the ages of the brown dwarfs are few Myr (triangles) and a broad age range for the field objects (triangles). The arrows shows the expected spin-up due to gravitational contraction. From Zhou et al. (2016).




Fig. 5.— The comparison between Earth, Jupiter, and stars shows that the magnetic energy density (in the dynamo) strongly correlates with a function of density and bolometric flux (here both in units of J m^{-3}). The bar lengths show estimated uncertainty rather than formal error. The stellar field is enlarged in the inset. Brown and grey ellipses indicate predicted locations of a brown dwarf with 1,500 K surface temperature and an extrasolar planet with seven Jupiter masses, respectively. From Christensen et al. (2009).




Fig. 6.— Predicted confidence regions for planet’s spin axis, from hypothetical single- and dual-epoch observations. Observing a planet at just a few orbital phases can significantly constrain both its obliquity and axial orientation. Obliquity is plotted radially: the centre is $= 0^\circ$ and the edge is $= 90^\circ$. The azimuthal angle represents the planet’s solstice phase. The green circles are the true planet spin axis, while the dark dashed lines and square show idealized constraints assuming perfect knowledge of the orbital geometry and kernel (i.e. no uncertainties). The upper left-hand and centre panels describe planet Q at phase angles 120° and 240° , respectively, while the lower left-hand panel incorporates both phases. For the colored regions, 10° uncertainty is assumed on each kernel width, inclination, and orbital phase, while 20° uncertainty is assumed on the change in dominant colatitude. Regions up to 3σ are shown, where darker bands are more likely. From Schwartz et al. (2016).




Fig. 7.— Albedo and its variations as a function of wavelengths for Solar System bodies with minor or no atmosphere. From Fujii et al. (2014).

5.2. B2: Which rocky planets have liquid water on their surfaces? Which planets have continents and oceans?

Relevance: Water is not a biosignature itself, but the presence of liquid water is required for life as we know it. Liquid water is not the only factor required for a planet to sustain life, but it is arguably the most important one. Thus, liquid water is a *habitability signature*. Establishing which habitable zone planets have liquid water on their surfaces provides an important context for EXOPAG SAG16, which focuses on biosignatures, but will rely on SAG15 for habitability signatures and characterization of habitable planets.

Our understanding of the distribution of water is surprisingly limited even for the case of Earth, and very incomplete for exo-earths: Currently, water detections (direct and indirect) in extrasolar systems are limited to protoplanetary disks (e.g., Carr & Najita 2008; Salyk et al. 2008), the atmospheres of hot jupiters and hot neptunes (e.g., Fraine et al. 2014), and in disks around white dwarfs fed by tidally disrupted minor bodies (Farihi 2016); however, no direct or indirect observations exist of water in extrasolar habitable zone Earth-like planets or even in super-Earths.

Simulations of exo-earth observations have been used to demonstrate that rotational phase mapping (time-resolved observations of hemisphere-integrated reflected light from the planet) can reveal the types and distribution of surfaces. Equipped with additional data on the color/spectra of the features and the physical conditions on the planetary surface may be used to identify surface features such as oceans and continents.

In the following we will discuss two different pathways for identifying liquid water on Earth-like habitable zone planets: 1) via the detection of oceans; and 2) via the detection of water clouds.

5.2.1. Detecting Oceans

The traditional habitable zone (HZ) is defined in terms of surface liquid water (Kasting et al. 1993). Three distinct methods have been proposed to search for large bodies of liquids (oceans) on the surface of a planet:

Polarization. For planets with low average ocean wind speeds ($\lesssim 2\text{m/s}$) oceans are smoother than other surface types (typically solids) and thus polarize light to a high degree (e.g. Williams & Gaidos 2008; Zugger et al. 2010, 2011). For idealized scenarios, the phase variations in polarization are significant, but in practice the effect of oceans is masked by

Rayleigh scattering, clouds, and aerosols. For planets with Earth-like average ocean wind speeds (≈ 10 m/s) the ocean surfaces (with the exception of the glint surface) will depolarize the reflected light (due to wind-induced ripples on the oceanic surface). Observations of polarized Earthshine, however, imply that rotational variations in polarized intensity may still be useful in detecting oceans (Sterzik et al. 2012).

Specular reflection The same smoothness that leads to polarization dictates that oceans are also able to specularly reflect light, especially at crescent phases (Williams & Gaidos 2008). The signal-to-noise requirements for phase variations are not as stringent as for rotational variations since the integration times can be much longer: weeks instead of hours. However, Robinson et al. (2010) showed that clouds not only mask underlying surfaces, but forward scattering by clouds mimics the glint signal at crescent phases, while atmospheric absorption and Rayleigh scattering mask the glint signature. They proposed using near-infrared opacity windows to search for glint, but this would only be possible if the effects of clouds could be accurately modeled for exoplanets. Moreover, Cowan et al. (2012a) showed that crescent phases probe the least-illuminated and hence coldest regions of a planet regardless of obliquity. Insofar as these planets have ice and snow in their coldest latitudes, then this latitude-albedo effect acts as false positive for ocean glint.

Rotational Color Variability: Although the faces of extrasolar planets will not be spatially resolved in the foreseeable future, their rotational and orbital motions produce detectable changes in color and brightness. Ford et al. (2001) used simulations of Earth to show that the changing colors of its disk-integrated reflected light encode information about continents, oceans, and clouds. The inverse problem — inferring the surface geography of a planet based on time-resolved photometry — is much more daunting than the forward problem.

Much progress has been made on the *exo-cartography* inverse problem since the seminal work of Ford et al. (2001). The rotational color variations of a planet can be used to infer the number, reflectance spectra, surface area, and longitudinal locations of major surface types (Fujii et al. 2010, 2011; Cowan et al. 2009, 2011; Cowan & Strait 2013). Meanwhile, the rotational and orbital color variations of an unresolved planet can be analyzed to create a 2-dimensional multi-color map equivalently a 2D map of known surfaces (Fujii et al. 2010; Kawahara & Fujii 2011, 2010; Fujii & Kawahara 2012).

5.2.2. *Liquid Water Clouds*

Additional methods may be used to deduce the probable presence of liquid water on the surface of a potentially habitable planet without directly or indirectly detecting an ocean. The presence of liquid water on the surface of an exoplanet can be indirectly inferred by the presence of liquid water clouds in the exoplanetary atmosphere. With the help of spectroscopy astronomers have detected signs of water vapor on a number of giant exoplanets and brown dwarfs and even water ice clouds on a brown dwarf (e.g., Skemer et al. 2016; Iyer et al. 2016; Brogi et al. 2014; Fraine et al. 2014).

Identifying clouds made of liquid water droplets (and not water ice) using polarization. On Earth, both liquid water and water ice clouds exist because liquid water is present on the surface. On an exoplanet, though, detection of water ice clouds could only be reliably be interpreted as a signature of surface liquid water if the surface temperature were independently known to be above freezing. Liquid water cloud detection is less likely to be a false positive for surface liquid water, although it could be in the presence of near-surface temperature inversions as may occur near the terminators of synchronously rotating planets. The detection of liquid water clouds can also be achieved with the help of broadband polarimetry. The state of polarization of starlight reflected by a planet is highly sensitive to the composition and structure of the planetary atmosphere. Observations of planets of our Solar system show that polarization is a powerful tool in the characterization of the micro- and macro- physical properties of clouds in planetary atmospheres (e.g., Hansen & Travis 1974; Mishchenko et al. 2010). Simulations of the polarization signal of terrestrial and gaseous exoplanets indicate that polarization can also be a powerful tool for the characterization of exoplanet atmospheres (e.g., Seager et al. 2000; Stam 2008; Karalidi et al. 2011). An early example of the power of polarimetry in the characterization of clouds in an exoplanetary atmosphere, is the retrieval of the cloud top pressure, and composition and size distribution of cloud droplets in the upper Venusian atmosphere using ground-based, unresolved observations of Venus by Hansen & Hovenier (1974).

The identification of the state of water clouds on Earth is routinely done with the help of polarization (Parol et al. 1995; Goloub et al. 2000). The (highly polarized) primary rainbow is a direct indication of the existence of liquid water clouds in a planetary atmosphere. Bailey (2007) was the first to suggest the use of the primary rainbow to detect liquid water clouds on exoplanets. Karalidi et al. (2011) and Karalidi et al. (2012) presented numerical simulations of broadband spectra of planets covered by a cloud deck and patchy liquid water clouds respectively, and showed that the rainbow is a robust tool for the detection of liquid water clouds in exoplanetary atmospheres.

Ice water clouds can interfere with the detection of liquid water clouds in the Earth’s

atmosphere. Ice clouds can produce highly polarized halos (rainbows), that could mask the primary rainbow of the liquid water clouds and the existence of the liquid water clouds altogether. However, Karalidi et al. (2012) showed that for a heterogeneous liquid and ice water cloud coverage like the Earth’s the primary rainbow of liquid water clouds will still be detectable. Even for extreme cases where optically thick ice clouds cover $\sim 50\%$ of the water clouds of an exo-Earth the primary rainbow will be detectable.

For an Earth-like planet orbiting at 1AU around a star at 10 pc the primary rainbow will appear between 0.03-0.044 arcsec from the parent star (phase angle of $\sim 30^\circ$ – $\sim 40^\circ$). To detect the primary rainbow we will need to observe the exoplanet with a spatial resolution of ~ 0.002 arcsec. **More discussion/references required on the feasibility of these proposed observations.**

Observational Requirements (draft)

Sample size:

Observations: 1) Polarimetry; 2) Time-resolved (rotational) multi-band photometry; 3) Time-resolved multi-epoch photometry

1. **Orbital semi-major axis** of a planet is critical as it may determine the presence of liquid water on the surface. How many visits per system are needed by a direct imaging mission to determine an accurate orbital distance?

2. **Presence of Greenhouses gases** and water vapor in the atmosphere: CO_2 and H_2O have strong features in the near-IR.

5.3. B3. What are the origins and composition of condensate clouds and hazes in ice/gas giants and how do these vary with system parameters?

Contributors: Daniel Apai, Anthony del Genio

All Solar System planets with an atmosphere also harbor condensate cloud and/or haze layers. Clouds and hazes influence the pressure-temperature structure of the atmosphere, its emission and transmission spectra, as well as its albedo. Particles or droplets that make up clouds primarily form through condensation and grow via further condensation and/or particle collisions. With grain sizes that may range from a micron to \sim millimeter, cloud particles/droplets have short settling time and are typical below the tropopause, where the dynamics of an atmosphere is most likely to saturate volatile constituents in regions of rising motion. Based on different extrapolations of clouds observed on Earth and on other Solar System planets, a range of cloud models have been proposed for giant exoplanets and brown dwarfs (for a review and comparison, see Helling et al. 2008). Haze particles (typically $< 0.1 - 1 \mu\text{m}$ in size) often form via photochemistry-driven (e.g., Venus and Titan) or charged-particles-driven chemical reactions in the upper atmospheres (< 1 bar); with long residence times these particles often introduce large optical depths to upper atmospheres. From an observational perspective clouds and hazes may also be used as tracers of atmospheric dynamics (circulation, mixing, turbulence). Presence of haze or cloud layers may also mask the presence of specific atmospheric absorbers even if present at large abundances at pressures higher than the particle layer.

Current Knowledge: Condensate clouds have been observed in brown dwarfs and in hot jupiters, over a very broad range of temperatures and pressures. High-altitude haze layers have been observed for transiting planets ranging from hot jupiters to super-earths and possibly for earth-sized planets, as well as for brown dwarfs. In the following we briefly summarize the key aspects of condensate clouds and haze layers and refer the reader to one of the many excellent in-depth reviews of the field for more complete summary.

Condensate Clouds: As the atmospheres of exoplanets encompass a very broad temperature range (~ 50 to $2,000$ K) these atmospheres are expected to harbor a large variety of condensates. For solar compositions the most important condensates include Ca-Ti-oxides, silicates, metallic iron, sulfides, CsCl and KCl, H_2O , NH_4HS , NH_3 , CH_4 (e.g., Lodders & Fegley 2002, for a recent review see Marley & Robinson e.g., 2015). Most of our current knowledge on cloud properties and compositions come from studies of Solar System planets (most importantly, Earth and Jupiter) and from the abundant samples of brown dwarfs. Water vapor and water ice clouds in Earth can be studied in-situ and via remote sensing; models developed to explain their behavior and properties are often used as a starting point for models of extrasolar clouds (Ackerman & Marley 2001), although it is likely that in

some exoplanet and brown dwarf atmospheres cloud formation and properties may be set by different processes (for a review of different cloud models see, e.g., Helling et al. 2008).

With over $\sim 3,000$ brown dwarfs known these objects provide a easy-to-study analogs of extrasolar giant planet atmospheres. Temperatures of known brown dwarfs range from ~ 250 K (below freezing point!) to above 2,300 K; an increasing number of known brown dwarfs have very low gravities and masses of only a few M_{Jup} , enabling the definition of samples essential for comparative parameter studies.

Comparative studies of brown dwarfs reveal the presence of silicate cloud layers through prominent infrared color-magnitude changes that occur through the M–L–T–Y spectral type sequence. The sequence itself is primarily set by the presence and absence of prominent gas-phase absorbers and not directly by the presence/absence of clouds (e.g., Burgasser et al. 2006; Cushing et al. 2006; Kirkpatrick et al. 2012); however, there is a strong correlation between the spectral type and colors of a given object (e.g., Burgasser et al. 2002; Burrows et al. 2006; Saumon & Marley 2008; Dupuy & Liu 2012). The general and oversimplified picture that emerged suggests that while the hottest (M-type) brown dwarfs are condensate cloud free, with temperatures sinking below $\sim 1,800$ K the atmospheres of L-type brown dwarfs are characterized by thick silicate clouds (resulting in red near-infrared colors between $1\text{--}3\ \mu\text{m}$); at even lower temperatures ($T < 1,300$ K) a transition to silicate cloud-free atmospheres is envisioned. Correspondingly, cool T-type brown dwarfs have blue near-infrared colors (dominated by scattering by gas molecules rather than particles), consistent with the lack of thick clouds in their upper atmospheres (see Figure 9). At even lower temperatures, within the Y spectral type, less refractory and less abundant species, including water ice, are expected to condense out and form clouds (e.g., Morley et al. 2014).

Although there is ample evidence supporting the overall picture described above, it is also clear that the above picture fails to capture the real complexity of cloud properties and atmospheric chemistry in brown dwarfs. Outstanding questions include the large dispersion in color along the L–T sequence (e.g., Burgasser et al. 2008); the unusually red colors of many of the very young brown dwarfs (and a few intermediate-age ones), a likely sign of unusually dusty upper atmospheres (e.g., Allers & Liu 2013; Liu et al. 2013). Furthermore, the first detections of water ice clouds has been reported in a Y-dwarf with an effective temperature of only ~ 250 K (Faherty et al. 2014; Skemer et al. 2016), enlarging the temperature range over which cloud models can be tested.

Recently, time-resolved high-precision observations (photometric and spectroscopic light curves) enabled the comparative studies of different cloud layers within the same objects, breaking the degeneracy between the effects of the multiple atmospheric parameters that may vary between any two brown dwarf (age, composition, temperature, surface gravity, vertical




Fig. 8.— Condensate clouds predicted for the upper atmospheres of giant planets of different temperature. By D. Apai, after Lodders (2003).

mixing, cloud structure). Space-based (HST and Spitzer) studies with sub-percent photometric precision found that most, if not all, brown dwarfs have heterogeneous (patchy) cloud cover (Buenzli et al. 2014; Metchev et al. 2015); ground-based surveys found that the highest amplitude brown dwarfs are at the L/T transition Radigan et al. (2014). Time-resolved spectroscopy of L/T transition dwarfs showed that the spectroscopic variations emerge from the atmospheres characterized by a mixture of warm thin cloud / cooler thick cloud patches, and *not* by clouds and cloud holes (Apai et al. 2013; Buenzli et al. 2015). Simultaneous HST (1.1-1.7 μm) and HST–Spitzer (1.1-1.7 μm and [3.6] or [4.5]) observations of clouds in L, L/T, and T-type brown dwarfs revealed pressure-dependent (vertical) structures with characteristic patterns for objects of different spectral types (Buenzli et al. 2012; Apai et al. 2013; Yang et al. 2015). Most recently, planetary-mass brown dwarfs and companions have also been accessible to rotational modulation studies, providing an opportunity to explore cloud properties as a function of gravity (Biller et al. 2015; Zhou et al. 2016; Leggett et al. 2016).

Directly imaged exoplanets and planetary-mass companions cover a spectral type range from early L to mid-T. These objects differ from old, high-gravity brown dwarfs both in the fine structure of their spectra (Barman et al. 2011a,b; Skemer et al. 2012) and, often, in their broad-band colors (see, e.g., Fig. 9), but show some strong similarities to some young brown dwarfs (e.g. Allers & Liu 2013; Faherty et al. 2013, 2016). From the small sample of directly imaged exoplanets it appears that early L-type exoplanets have colors similar to brown dwarfs with matching spectral types (), late L and L/T-type exoplanets are often much redder and fainter than brown dwarfs with matching spectral types (e.g., Chauvin




Fig. 9.— Condensate clouds have a fundamental impact on the positions of brown dwarfs and directly imaged exoplanets on the near-infrared color-magnitude diagram. Along the L-type sequence (red) silicate clouds in the upper atmosphere become thicker. The cooler T-dwarfs are bluer because the silicate clouds are below the visible upper atmosphere. Figure from Wagner et al. (2016), which is in part based on the parallax database by Dupuy & Liu (2012).

et al. 2005; Marois et al. 2008), but the coolest T-type exoplanets appear to have colors consistent with those of T-type brown dwarfs (Macintosh et al. 2015; Wagner et al. 2016). This pattern, if verified, would argue for a difference in cloud properties (most significant in late-L and L/T transition objects) between the higher gravity brown dwarfs and the low-gravity exoplanets.

Clouds have also been studied in hot jupiters via transmission and emission spectroscopy, spectral phase mapping, and in reflected light. Observations from the Kepler space telescope (dominated by reflected light) argued for a large-amplitude, heterogeneous silicate cloud cover (e.g., Demory et al. 2013) that avoids the cold trap in the night side of the planet. (Fortney et al. 2008) proposed that the presence/absence of silicate clouds in hot jupiters should follow the general sequence observed in brown dwarfs. Although optical-near infrared HST transmission spectra argued for the presence of cloud decks in some hot jupiters (Gibson




Fig. 10.— Gemini/GNIRS spectrum of the ~ 250 K Y-dwarfs WISE0855 shows a series of absorption features attributable to water vapor, muted by clouds (likely water ice). From Skemer et al. (2016).

et al. e.g., 2013; Sing et al. e.g., 2015), no clear trend (in terms of presence/absence of clouds) emerged from a homogeneous survey of hot jupiters (Sing et al. 2016). In contrast, Stevenson (2016) suggests that clouds in hot jupiter atmospheres are restricted to regions in the surface gravity/temperature plane. It is likely that the presence of silicate clouds in the regions probed by transmission (terminator) and emission spectroscopy (dayside) strongly depends on the day-night temperature difference (e.g., Rauscher & Menou 2013), atmospheric circulation (see Question C1, and e.g., Showman et al. 2009, 2015), and the importance of potential cold traps (Parmentier et al. 2013).

Kepler-measured planet phase curves contain contribution of both reflected light and planetary thermal emission. Distinctive phase dependency of the two components may allow them to be separated (Hu et al. 2015), and the reflective component is directly related to the distribution of clouds on the planet. This method has been applied to three hot Jupiters, and they all appear to have heterogeneous silicate clouds (Demory et al. 2013; Shporer & Hu 2015). Detailed models involving cloud condensation and general circulation suggest that such heterogeneous clouds are indeed common on hot Jupiters, and the cloud-forming material differs under different temperature regimes (Parmentier et al. 2016). This knowledge is relevant for direct imaging because (1) it proves that at least some exoplanets have highly reflective clouds and therefore high albedo, and (2) it calls for considering inhomogeneous cloud coverage when interpreting spectra from direct-imaging observations.

Hazes – with particles less than $0.1 \mu\text{m}$ – have been argued for in a few objects where

the lack of near-infrared absorption features (commonly water) necessitates that the absorption features are muted by high-altitude particles (unless the upper atmospheres of some transiting planets are extremely dry, see Madhusudhan et al. 2014). Such strong reduction in water absorption features was seen in the warm sub-neptune GJ1214b (Kreidberg et al. 2014) and atmospheric models argued for the presence of very small particles at low pressures (~ 1 mbar), consistent with photochemical hazes but not with condensate clouds (Morley et al. 2015). In an ongoing study, the transmission spectra of moderate earth-sized planets around a very low-mass red dwarf also appear to be flat, perhaps also influenced by small particles lofted to low pressures (de Wit et al. 2016, in prep.). The detection of hazes (based on a very similar water absorption-based evidence) argues for the presence of some haze in L-type brown dwarfs, in spite of the lack of a host star, which argues for non-photochemical haze production (Yang et al. 2015), possibly driven by charged particles accelerated by the brown dwarf’s magnetic field.

Solar System Gas Giants as exoplanet analog observations: Overlapping Kepler photometry and Hubble Space Telescope images of Neptune have shown complex time-varying signal whose frequency analysis revealed not only the fundamental rotation rate, but also the level of differential rotation of major mid-latitude cloud features (Simon et al. 2016). Quasi-continuous 20-hour-long two-band optical imaging of Jupiter with the Hubble Space Telescope provided simultaneous high-precision photometry and high-fidelity and high-resolution images (Karalidi et al. 2015). These authors showed that MCMC-based lightcurve modeling can correctly retrieve the position, size, and surface brightness of the dominant features in the lightcurve, such as the Great Red Spot, even from a single rotation.

Sub-questions

The study of extrasolar cloud layers is novel and we may not be in the position yet to identify the right set of key questions to ask. Nevertheless, the following list attempts to capture the most important uncertainties of our current models of clouds.

- What are vertical structures of single and multi-layer clouds formed from different condensates?
- What are the grain size distributions and compositions (single-species or compound grains) in the clouds?
- Under which conditions do photochemically- and charged particle-driven haze layers form? How complex can chemistry get in haze layers?
- How do condensate clouds form and evolve as a function of fundamental atmospheric parameters?

Observational Requirements (draft)

Sample size:

Observations:

Questions to SAG15:

1) How challenging are the very different methods described here relative to each other?

2) How are they best carried out?

Kasper: ELT: A Jupiter analogue (1Rj, 5AU) will just be observable ($c \sim 1e-9$). Neptune / Uranus analogues cannot be seen with an ELT.

5.3.1. Science Value of Independently Measured Planet Masses and Radii

5.4. B4. How do photochemistry, transport chemistry, surface chemistry, and mantle outgassing effect the composition and chemical processes in terrestrial planet atmospheres (both habitable and non-habitable)?

The composition of exoplanetary atmospheres is one of the central questions of exoplanet characterization. Atmospheric composition is influenced by a multitude of factors (both initial and boundary conditions and processes) and, therefore, can provide valuable insights into the formation and evolution of each planet as well as on its present-day status. For example, characterizing the atmospheric composition of habitable zone planets is essential for determining whether they are, in fact, habitable planets — in other words, that surface conditions allow the presence of liquid water (see also §5.2).

Direct imaging missions are expected to image a diverse range of planets both in mass (from sub-earths to super-jupiter), in temperature (<100 K to >500 K), and in composition (H-rich, CO_2 -dominated, atmosphereless, etc.). In the initial reflected light images, identifying the type of planets in a system may be very difficult (e.g., a small but high-albedo planet may look identical to a large but low-albedo planet; or — even if the albedos are similar — a partially illuminated (crescent-phase) giant planet may be very similar to a full disk of a slightly hazy super-earth). It is therefore imperative for direct imaging missions to include some level of planetary characterization as part of a discovery survey.

A multitude of excellent and up-to-date reviews are available on exoplanet atmospheric composition, based on observational evidence (Solar System planets, brown dwarfs, and exoplanets), and on theoretical predictions for the range of possible and expected compositions; therefore, we focus on the questions most salient for direct imaging missions.

Example sub-questions

- a) What are the major and minor constituents of the atmospheres of rocky planets?
- b) How do the compositions of rocky planet atmospheres vary as a function of mass, bulk composition, and irradiation?
- c) How strongly does mantle outgassing affect rocky planet atmospheres?
- d) Which planets show evidence for primordial atmospheres? item e) How are planetary atmospheres impacted by stellar high-energy radiation and stellar wind?

Example Science Cases and Observations

Thorough exploration of the possible compositional classes for warm super-earth/neptune atmospheres in argues for at least six classes (see Fig. 11): i) Water-rich atmospheres; ii) co-existing water and hydrocarbon-dominated atmospheres; iii) hydrocarbon-rich atmospheres; iv) oxygen-rich atmospheres; v) CO/CO₂-dominated atmospheres; and, vi) H₂/He-rich atmospheres. The photochemistry in these atmosphere types has been explored, for example, in Hu & Seager (2014). Many smaller terrestrial planets are likely to have CO₂- or N₂-dominated atmospheres, based on solar system experience, with significant amounts of sulfur-bearing gases if volcanic activity is present (Hu et al. 2012b, 2013).




Fig. 11.— The types of thick atmospheres possible on Super-Earths and mini Neptunes, based on the extensive exploration of chemical reaction networks. For atmospheres *not* dominated by H₂, different atmosphere classes emerge as a function of the relative abundances of C, O, and H. From Hu & Seager (2014).

5.4.1. Science Value of Independently Measured Planet Masses and Radii

6. Exoplanetary Processes

6.1. C1. What processes/properties set the modes of atmospheric circulation and heat transport in exoplanets and how do these vary with system parameters?

Authors: Daniel Apai, Nick Cowan, Ravi Kopparapu, Anthony del Genio

Atmospheric circulation plays a key role in redistribution of the energy in exoplanet atmospheres. Depending on typical wind speeds, rotational velocity, insolation, latent heat released during condensation, and other system parameters different atmospheric circulation regimes are expected on planets that can be studied with direct imaging missions. For potentially habitable exoplanets atmospheric circulation will determine the day-night heat differential and the equator-pole temperature difference. Understanding the presence and size of Hadley cells can also provide important insights into how water vapor (or other condensibles) may be distributed in habitable planets.




Fig. 12.— Depending on the relative importance of rotational speed, wind speed, and vertical heat transport, simple models predict two different regimes of circulation for giant planets: vortex-dominated (left) and jet-dominated (right). From Zhang & Showman (2014).

Understanding atmospheric circulation in habitable exoplanets is an important component in establishing a correct climate model for them. As of now, atmospheric circulation has modeled in the Solar System planets and a small sample of brown dwarfs, hot jupiters and lower-mass exoplanets (see Figs 14 and 15, Yang et al. 2013; Leconte et al. 2013; Abe et al.




Fig. 13.— Dependence of the atmospheric circulation on rotation rate. Panel (a) shows a slowly-rotating, highly irradiated hot-Jupiter planet with strong day-night temperature difference and a strong eastward equatorial super-rotating jet. Panel (b) shows rapidly rotating warm Jupiters that are weakly irradiated. These planets develop eddy driven zonal jets that peak at mid-latitudes rather than at the equator. From Showman et al. (2015).

2011; Wordsworth et al. 2011; Zhang & Showman 2014; Kopparapu et al. 2016; Kataria et al. 2014; Wolf & Toon 2014). The nature of the atmospheric dynamics depends on the thickness of the planet’s atmosphere, its rotation rate, the distance of the planet from the star and several other factors. A more comprehensive study of different atmospheric circulation regimes of exoplanets still lacks, but important steps have been taken for rocky exoplanets in a simplified general circulation model by (Kaspi & Showman 2015).

The atmospheric circulation patterns of planetary atmospheres can be characterized broadly from the planetary rotation rate; Earth exhibits three major circulation cells, while planets with a more rapid rotation rate and/or larger radii (such as gas giants) show five or more circulation bands (Williams & Holloway 1982). Knowledge of an exoplanet’s rotation rate would provide a strong constraint on the large-scale dynamical features that should occur, given the planet’s orbital distance from its host star (Merlis & Schneider 2010).

Different circulation regimes can exist in the atmospheres of extrasolar planets depending upon the incident flux and rotation rate of planet. For example, (Showman et al. 2015) showed that the canonical hot-Jupiter regime (0.03 – 0.05 AU), with a large day-night temperature gradient and a fast east ward equatorial jet, transitions at lower stellar fluxes (~ 1 AU) and/or faster rotation to a regime with small longitudinal temperature variations and peak wind speeds occurring in zonal jets at mid- to high latitudes.

Furthermore, at a given stellar flux, a greater than factor of two in rotation rate difference between synchronous and non-synchronous causes can potentially be discerned in the light curves of hot-Jupiters, providing a way to identify regime transition from highly




Fig. 14.— Moist/water-rich atmosphere simulations from (Wolf & Toon 2015). The four panels indicate the amount of cloud water content on a planet at different insolation levels (or, alternately, how close to an inner edge of the HZ a planet is located). From left to right, the solar insolation varies: S_0 (current Earth insolation), 10% of S_0 , 12.5% S_0 and 21% of S_0 . This is for an Earth-size planet around a Sun-like star.

irradiated to weakly irradiated planets.

Observational characteristics such as variation in thermal emission from orbital phase curves, and net Doppler-shift obtained from high resolution spectra taken during the transit can, in principle, provide a means to constrain the rotation rate for some hot-Jupiter planets (Rauscher & Kempton 2014). Although these techniques may not be individually suited to distinguish the rotation rates, the combination of these two techniques may show observable differences with rotation rate.

Planets in and around the habitable zone (HZ) of low-mass stars are expected to be in synchronous rotation, though thermal tides can cause asynchronicity on some planets (Leconte et al. 2015). Such planets can further be classified as *slow-rotators* (where the Rossby deformation radius is equal or greater than planetary radius) and *fast-rotators* (where the Rossby deformation radius is less than planetary radius). Planets in synchronous orbits that are also slow-rotators may develop a shielding cloud presence beneath the substellar point, which can increase the inner habitable limits of the planet (Yang et al. 2013). However, rapidly-rotating planets tend to smear out this cloud deck, which limits much of this shielding effect (Kopparapu et al. 2016).

On Earth, the mean meridional circulation, or Hadley circulation, is responsible for the poleward transport of energy at low latitudes; however, on synchronously rotating planets, the Hadley circulation provides an incomplete diagnosis of energy transport because the Hadley circulation itself changes direction between the hemisphere eastward and westward of the substellar point (Haqq-Misra & Kopparapu 2015) and a significant day-night circulation develops when the radiative time scale is shorter than the length of the solar day (Way et al.




Fig. 15.— Temperature and horizontal wind vectors at the surface, 0.5 bar, and 0.1 bar levels for an Earth-mass planet in a slow-rotating regime near the inner edge of the habitable zone around a K-dwarf. Slowly rotating planets develop sub-stellar clouds that increase the albedo of the planet. Inflow along the equator and from the poles into the substellar point at the center is also shown. From Kopparapu et al. (2016).

2016). Rather than the Hadley circulation, the mean zonal circulation (or Walker circulation) provides a better metric for synchronous rotators to examine the efficacy of heat transport between the substellar and antistellar points. For slow rotators, the Walker circulation reaches to the night side of the planet, but for rapid rotators, the Walker circulation by itself is limited in longitudinal extent. In such cases, a cross-polar circulation also provides energy transport between the day and night side to keep the atmosphere from freezing-out or collapsing (Joshi et al. 1997; Haqq-Misra & Kopparapu 2015).

Recent three-dimensional climate modeling studies of Earth-like planets predict that rapidly rotating planets undergo a sharp transition between temperate and moist greenhouse climate states (Wolf & Toon 2015; Popp et al. 2016). Wolf & Toon (2015) argue that this transition is associated with a fundamental change to the radiative-convective state of the

atmosphere. When the mean surface temperature approaches ~ 330 K, the lower atmosphere becomes opaque to infrared and thermal radiation due to increasing water vapor mixing ratios. The lower atmosphere heats due to solar absorption in the near-IR. Simultaneously, the lower atmosphere cannot efficiently cool to space due to the closing of the 8-13 μm water vapor window region. Combined, this results in a net positive radiative heating rates in the near surface layers, creating a ubiquitous temperature inversion across the planet. The inversion suppresses boundary layer convection, reducing clouds and the planetary albedo at the climatic transition. As climate warms further, the low atmosphere becomes increasingly hot and dry (i.e., low relative humidity), but upper atmosphere water vapor mixing ratios become large and a zonally uniform, albeit patchy, cloud deck develops.

Figure 14 shows the evolution of the zonal mean cloud water content (kg m^{-3}) for an Earth-like planet under increasing stellar fluxes, varying from the present day Earth insolation up to a 21% increase. For the present day Earth climate (Fig. 14, leftmost panel), clouds are confined to pressures greater than ~ 200 mb, with the thickest clouds located at mid-latitudes. For moist greenhouse atmospheres, the lower atmosphere becomes cloud free, while the primary cloud deck becomes zonally uniform and is pushed higher in the atmosphere. For an Earth-like planet with a mean surface temperature of ~ 363 K, the cloud water peaks near ~ 50 mb (Fig. 14, rightmost panel). Clouds are well known to obscure exoplanetary spectra due to their significant broadband opacity. Thus we may be able to differentiate habitable Earth-like atmospheres from moist greenhouse atmospheres, based on the pressure level of the primary cloud deck. However, note that an Earth-like planet at ~ 363 K, would have moist stratosphere ($\sim 6 \times 10^{-2}$ H_2O mixing ratio at 0.2 mb), and thus would be expected to lose an Earth ocean of water to space within several hundred million years. Moist and runaway greenhouse atmospheres are thus transient phenomena.

Interestingly, Kopparapu et al. (2016) found that the above described radiative-convective transition also occurs on slow and synchronously rotating Earth-like planets, which are expected around low mass stars. While rapidly rotating planets can maintain climatological stability beyond this transition due to cloud adjustments in the upper atmosphere, this transition is catastrophic for planets located near the inner edge of the habitable zone around low mass stars. As noted above, Synchronously rotating planets are effectively shielded from the host star by thick convectively produced clouds located around the substellar point. These planets can remain habitable despite incident stellar fluxes up to twice that of the present day Earth (Yang et al. 2014; Kopparapu et al. 2016). However, the radiative-convective transition and subsequent onset of the near surface inversion stabilizes the substellar atmosphere, and thus the convective cloud deck rapidly dissipates. Even a small dent in this substellar cloud shield then lets in a tremendous amount of solar radiation, destabilizing climate towards an immediate thermal runaway.

Questions to SAG15:

To what level can the atmospheric circulation be constrained for different types of planets?

What hypotheses / toy circulation models should be tested for gas giants?

What hypotheses / toy circulation models should be tested for habitable super-earths / earths?

What data type and cadence is required or best suited for characterizing circulation?

How does the atmospheric circulation in tidally locked planets around M-dwarf stars affect habitability?

Where is the transition region from slow to rapid rotators in tidal-locked planets around low-mass stars?

Observational Requirements (draft)

Sample size:

Observations: Multi-epoch near-IR spectral observations at moderate to high resolutions

6.2. C2. What are the key evolutionary pathways for rocky planets and what first-order processes dominate these?

Contributors: Nick Cowan, Daniel Apai, Renyu Hu

The two earth-sized rocky planets in the Solar System, Earth and Venus, likely started with very similar initial mass, orbit, and composition, but their evolutionary paths have strongly diverged. Mars, although substantially different in its mass and orbit, has again followed a different evolutionary trajectory, even though it is thought that surface conditions on early Mars, at least temporarily or episodically, may have sustained wide-spread aqueous activity on the the surface, perhaps resembling the early Earth. With the large number of rocky planets that may be observable with a capable future direct imaging mission, the range of evolutionary histories could be explored.

The question naturally emerges: What key evolutionary pathways exist for rocky planets and what factors determine which of these pathways a given planet will follow?

Attractors and Divergence in the Phase Space of Rocky Planet Evolution: It is reasonable to describe the momentary state of a given rocky planets with a set of n fundamental parameters and explore the evolution of the planet in this n -dimensional phase space. Each planet’s history and future evolution is thought of as a trajectory. Fundamental parameters could include, but are not limited to, planet mass, radius, atmospheric pressure scale height, orbital parameters, atmospheric composition, rotation rate, magnetic field strength, etc. Which trajectory a planet follows will depend not only in its momentary location in the phase space, but also by the effect of a set of feedback loops (both positive and negative) as well as on a few environmental variables (e.g., stellar luminosity and incident optical and UV flux).

When describing planet evolution in such a manner, several obvious questions are identified: 1) *How sensitive are the trajectories to initial parameters and/or perturbations to the system?* 2) *What is the importance of a planet’s past, e.g., which volumes of the phase space are uni-directional (e.g., irreversible water loss)?* 3) *Are there preferred evolutionary end-states (attractors) or is the surface defined by coeval planets smooth?* 4) *What is the importance of quasi-monotonic evolution driven by a small number processes vs. random walk driven by a multitude of competing processes?*

Exploring the past history and current state of rocky planets allows the system-level study of rocky planet evolution and will be essential for understanding the occurrence rate of truly earth-like planets and to place the physical processes that drive planet evolution on Earth to the broader context of exo-earths.

6.2.1. Science Value of Independently Measured Planet Masses and Radii

Is the science goal achievable without precise mass measurements? Yes, a medium or large sample of rocky exoplanets for which most of the other key parameters are known would likely suffice to establish the topology of the phase space.

Would the science goal benefit greatly from precise mass measurements? Yes, precise mass measurements would significantly contribute to the understanding of the planets' properties. In case the phase space is highly complex and its projection to a lower-dimensional (observed) phase space does not allow the identification of the key processes that drive the evolution, expanding the projected phase space by a new dimension (mass) may break the degeneracy between different processes that lead to similar evolutionary outcomes.

Observational Requirements (draft)

Sample size: probably large samples are required (>50–100)

Observations: characterization of the planets: atmosphere pressure and composition, orbital parameters, bulk composition, surface temperature estimate, stellar parameters and past evolution;

Comments: To explore: Toy model for testing hypothesis of smooth distribution vs. attractors

6.3. C3. What types/which planets have active geological activity, interior processes, and/or continent-forming/resurfacing processes?

Contributors: Stephen Kane, Daniel Apai, Nick Cowan

Planetary interior processes and geological activity play an important role in coupling Earth’s atmosphere to its crust and providing a long-term stabilizer for Earth’s climate. The source of Earth’s atmosphere and volatiles are mostly products of outgassing after the loss of the primary atmosphere. Developing reliable climate models to determine the habitability of potentially habitable planets will likely require assumptions about the geological activity and the level of coupling between the planet’s crust and atmosphere (e.g., Abbot et al. 2012; Foley & Driscoll 2016). However, interior processes are obviously very difficult to probe via low signal-to-noise and spatially unresolved remote sensing.

The influence of geological activity on planetary climate is most clearly understood for the case of Earth. On geologic timescales, continental crust production participates in the stabilization of the Earth’s climate through its role in carbonate weathering feedback. Chemical weathering of silicate minerals on land in the presence of water causes the slow removal of CO_2 from the atmosphere, which is eventually deposited on the ocean floor as carbonate compounds. Without the continual re-injection of new CO_2 by volcanoes, the atmospheric stock of CO_2 would be slowly depleted. However, the rate of CO_2 removal by silicate weathering is temperature dependent, so that in the presence of a steady source of volcanic CO_2 , weathering interacts with the greenhouse properties of CO_2 to produce a negative feedback on planetary temperature. This interaction, whereby warmer conditions lead to increased drawdown of CO_2 and a consequent weakening of the greenhouse effect (and vice versa), is believed to play an important role in stabilizing planetary temperatures in the presence of a main-sequence star which is increasing in luminosity over Ga timescales. It is because of this process that it has been argued that volcanism and geological activity are necessary conditions for sustained life on a planet.

Current Knowledge: Two methods have been proposed to detect geological activity on a rocky exoplanet. First, Kaltenegger & Sasselov (2010) suggested that volcanic emission of SO_2 can be detected remotely. However, it has been found that the volcanic sulfur emission would most likely lead to formation of sulfur and/or sulfate aerosols in the atmosphere, leading to muted transmission and thermal emission spectral features Hu et al. (2013). The sulfur-bearing aerosols may be detected via direct imaging, and indicate volcanic activity on the planet. Second, Hu et al. (2012a) suggest that fresh volcanic surfaces and surfaces solidified from a magma ocean have prominent spectral features at 1 micron and 2 micron, produced by Si-O bonds in mineral lattices. Surfaces aged by either space or aqueous weathering do not have these features. Therefore, concern spectral features can

imply recent volcanic activities on a rocky exoplanet.

Studies of terrestrial climate and volcanism focus primarily on the effects of volcanism on surface temperature, which we are unlikely to be able to estimate for most exoplanets. However, volcanically forced anomalies in surface temperature are coupled to anomalies in emission temperature, which can be targeted for follow-up observations. Thus, if volcanism can be identified on an exoplanet it may represent the most promising method for estimation of climate sensitivity outside of the Solar System. **citations?**

The distinctive effect of volcanic eruptions on the transmissivity of atmospheres is related to the force of their explosions. Typically, processes on Earth that produce aerosols in the atmosphere affect only the troposphere. Aerosols are quickly washed out of the troposphere by rain, and thus a sustained impact on atmospheric transmissivity requires a near-continual source of the aerosol or its precursor gas. Many small eruptions don't reach the stratosphere, however the largest explosive volcanic material can, in contrast, inject SO_2 directly into the stratosphere, where it reacts to form sulphate aerosols (e.g., Kaltenegger & Sasselo 2010).

Because the stratosphere is very dry and the particle sizes are small, these aerosols can persist in the stratosphere for several years, until they are removed by the natural overturning circulation of the stratosphere (Robock et al. 2007). Stratospheric air rises in the tropics and then migrates towards the pole where it sinks. Because of this, aerosols from tropical eruptions typically persist in the stratosphere for about two years, while aerosols from high-latitude volcanism persist for only one year (Robock et al. 2007; Tingley et al. 2014).

Previous work shows a link between exoplanet compositions and stellar compositions (e.g., Rogers & Seager 2010) such that stellar compositions can be used to approximate the composition of exoplanet interiors. Stars in exoplanetary systems show a wide variation in composition (Hinkel et al. 2014). In particular, some composition parameters with large variability such as Mg:Si ratios, are likely to have a first order effect on the minerals that compose exoplanetary interiors and thus the melting behavior, magma composition generated from these planetary mantles, and their volatile solubility. Certain compositional components, such as alkalis, have also been shown to greatly increase the H_2O solubility (e.g., Behrens & Zhang 2001; Larsen & Gardner 2004) in natural melts, and highlight the necessity of measuring volatile solubility behavior across a broad range of melt compositions. Magmatic volatile solubility is highly dependent on temperature, which also varies with mineralogy.

On Earth, in addition to the pressure- and compositional-dependence of volatile solubility in magmas, the explosivity of a given eruption is dependent on the overall volatile

concentration (dominated by H_2O and CO_2), magma supply rate, vent geometry, and source pressure of the magma body (e.g., Wilson 1980; Papale & Polacci 1999; Mason et al. 2004). The most explosive eruptions on Earth tend to be those at convergent plate boundaries where there are abundant volatiles involved in magma genesis sourced from the subducting plate, and some types of intraplate volcanism where interactions with reservoirs of volatiles in the crust produce highly explosive caldera eruptions. In addition, flood basalts and other volumetrically large outpourings of magma common in a planets early history may be a significant source of atmospheric volatiles (Black et al. 2012). As such, the lack of tectonics on exoplanets does not preclude extreme volcanism that may produce detectable signatures.

6.3.1. *Geological Activity and Plate Tectonics on Extrasolar Rocky Planets*

The terrestrial and venutian mantle convection, plate tectonics, and mantle outgassing are influenced by the initial bulk abundance of the planet and are particularly sensitive to the radioisotopic abundances; mantle outgassing and planetary evolution are particularly sensitive to the the modes of the tectonics (e.g., stagnant lid vs. plate tectonics), internal temperature distribution, and lid thickness (e.g., O’Neill et al. 2013). The extrapolation of models of planetary evolution and plate tectonics to extrasolar rocky planets is challenging. A particularly relevant question is how plate tectonics may operate in super-Earths: on one hand, the higher heat flux (due to their intrinsically higher mass-to-surface ratio) should lead to stronger mantle convection (e.g., Valencia et al. 2007; van Heck & Tackley 2011). On the other hand, based on a visco-elastic models of mantle convection and crust formation, O’Neill & Lenardic (2007) find that increasing the planet’s radius (and mass) will decrease the ratio of driving-to-resistive forces (see Fig. 16), which reduces the likelihood of mobile plate tectonics in super-Earths and argues for the stagnant lid (or episodic tectonics) in these planets.

Furthermore, for a given planet models also suggest time-dependence and sensitivity to initial conditions: the thermal state of the post-magma ocean mantle is a key parameter that determines the subsequent evolution of the planet (possibly but not necessarily through i) hot stagnant-lid, ii) plate tectonics, then to iii) cold stagnant lid regime). Depending on the planet’s transition from the magma ocean stage different evolutionary paths are possible and there may only be a limited time available for Earth-like plate tectonics (O’Neill et al. 2016).




Fig. 16.— Convection as a function of stellar radius and Byerlee-style pressure-dependent yield stress. The models include internal heating, a constant friction coefficient, and gravity matching the planetary mass. Larger radius results in greater buoyancy forces, but also increased fault strength due to increased pressure. Thus planets with larger radii again tend to be in an episodic or stagnant regime, depending on the absolute yield stress. From O’Neill & Lenardic (2007).

6.3.2. Observational Methods

While major geological processes usually unfold on timescales not accessible to long-range remote sensing, the *results* of these processes are detectable and, in some cases, may be unambiguously identifiable. For example, in the case of Earth the presence of multiple large land-masses and oceans (detectable via time-resolved observations, e.g., Cowan et al. 2009) reveals that a continent-forming process acts on timescales shorter than water-driven land erosion and provides a characteristic scale for the continental plates. Another Earth-based example is the accumulation of atmospheric absorbers characteristic of volcanic outgassing (e.g., SO_2 : Kaltenegger & Sasselov 2010). Other, non-Earth-like, planets may offer other detectable signatures of geological activity.

In the following we briefly discuss four representative possibilities:

- i) Continents and Oceans from Surface Maps
- ii) Atmospheric Absorbers from Volcanic Outgassing
- iii) Planetary-Scale Surface Mineralogy
- iv) Cloud formation as Tracer of Topography and Erosion

Continents and Oceans from Surface Maps: Simulated observations of Earth as an exoplanet demonstrate that with appropriate rotational- and orbital phase-resolved precision, multi-band photometric data can be used to identify the presence and one-dimensional and two-dimensional distribution of oceans and landmasses (see also Section 5.2.1, e.g., Cowan et al. e.g., 2009; Fujii et al. e.g., 2010; Fujii & Kawahara e.g., 2012). In a planet where large bodies of liquid water (an ocean) is present, a hydrological cycle is active, and land masses (continents) are detected, land erosion must arguably occur; the timescales for the erosion may be, to the first order, estimated based on terrestrial silicate weathering and erosion rates. The existence of the continents demonstrates that the time-scale of continent-formation is comparable or faster than their erosion. Based on a simplified model of water cycling and continent formation, Cowan & Abbot (2014) argues that continents and oceans may be common even among super-earths with high abundances of water. Such first-principle-based models may be combined with the scales of oceans and continents derived from observations to test whether active continent formation (e.g., plate tectonics) is required for a given planet.

Water Clouds as Tracer of Topography Similarly to the previous Pallé et al. (2008)

Planetary-scale Surface Geology Common mineral assemblages that make rocky planet surfaces have distinctive spectral features in the visible, near-infrared, and thermal infrared wavelengths. Broadband photometry of atmosphere-less rocky exoplanets can therefore tell their surface types (Hu et al. 2012). For example, water-altered silicate surfaces (e.g., clays) will produce narrow absorption bands at 1.8 and 2.3 micron owing to the OH incorporated in the solids. For another example, the location of the peak in the 7-10 micron band of a silicate rock tells its silica content, which can be used to distinguish primary versus secondary crust on a rocky planet. Fujii et al. (2014) used albedo-map generated lightcurves and, where available, observed photometric variations to explore the geologic features detectable on diverse Solar System bodies with minor or no atmospheres (Moon, Mercury, the Galilean moons, and Mars). The study included the evaluation of the light curves and the features that are detectable at wavelengths ranging from UV through visible to near-infrared wavelengths, and also explored the accuracy required to determine the orbital periods of these bodies.

Figure 7 provides an example for the wavelength-dependence of the rotational variability amplitudes in different bodies.

Amplitude variations at the level of 5–50% have been reported introduced by features of diverse nature (volcanism, space weathering, planetary weathering, impact excavation, tectonic deformation). In some cases data with the appropriate wavelength coverage can be used to identify some of these features or narrow down the possible origins.

Volcanic Outgassing

6.3.3. Complementary Datasets

We identify three complementary datasets that are critically important for modeling the interior and activity of extrasolar rocky planets:

- *Stellar abundances*: a proxy for the relative refractory elemental abundances that may be present in the planet; may be used to identify outlier systems in terms of elemental abundances
- *Composition of giant planets in the system*:
- *Stellar/system age*: to constrain the evolutionary state of the planets (heat flux and time available for volatile loss and resurfacing processes)
- *Mass and radius of the planet*: fundamental physical parameters with major impact on energy budget and force balances; and constrains the bulk composition

6.3.4. Science Value of Independently Measured Planet Masses and Radii: Very High

Exploring the planetary-scale geophysics of rocky planets will likely be among the most challenging aspects of characterizing extrasolar rocky planets. Yet, understanding the geophysics and interior activity of these planets may well turn out to be essential for correctly and robustly interpreting atmospheric biosignatures. The rocky planet’s mass is one of the most fundamental parameter that influences heat flux, pressure, and horizontal forces acting on the lithosphere. Given the sensitivity of plate tectonics models to planet mass, it is likely that determining the planet mass with a precision of $\sim 10\%$ is required for establishing a robust geophysical model.

7. Data Requirements

This section identifies the type and quality of data ideal or required to answer the individual science questions.

Appendix

A. SAG15 Charter

Future direct imaging missions may allow observations of flux density as a function of wavelength, polarization, time (orbital and rotational phases) for a broad variety of exoplanets ranging from rocky sub-earths through super-earths and neptunes to giant planets. With the daunting challenges to directly imaging exoplanets, most of the community’s attention is currently focused on how to reach the goal of exploring habitable planets or, more specifically, how to search for biosignatures.

Arguably, however, most of the exoplanet science from direct imaging missions will not come from biosignature searches in habitable earth-like planets, but from the studies of a much larger number of planets outside the habitable zone or from planets within the habitable zone that do not display biosignatures. These two groups of planets will provide an essential context for interpreting detections of possible biosignatures in habitable zone earth-sized planets.

However, while many of the broader science goals of exoplanet characterization are recognized, there has been no systematic assessment of the following two questions:

- 1) What are the most important science questions in exoplanet characterization apart from biosignature searches?
- 2) What type of data (spectra, polarization, photometry) with what quality (resolution, signal-to-noise, cadence) is required to answer these science questions?

We propose to form SAG15 to identify the key questions in exoplanet characterization and determine what observational data obtainable from direct imaging missions is necessary and sufficient to answer these.

The report developed by this SAG will explore high-level science questions on exoplanets ranging from gas giant planets through ice giants to rocky and sub-earth planets, and – in temperatures – from cold (~ 200 K) to hot ($\sim 2,000$ K). For each question we will study and describe the type and quality of the data required to answer it.

For example, the SAG15 could evaluate what observational data (minimum sample size, spectral resolution, wavelength coverage, and signal-to-noise) is required to test that different formation pathways in giant planets lead to different abundances (e.g. C/O ratios). Or the SAG15 could evaluate what photometric accuracy, bands, and cadence is required to identify

continents and oceans in a habitable zone Earth-sized or a super-earth planet. As another example, the SAG15 could evaluate what reflected light data is required to constrain the fundamental parameters of planets, e.g. size (distinguishing earth-sized planets from super-earths), temperature (cold/warm/hot), composition (rocky, icy, gaseous), etc.

SAG15 will not attempt to evaluate exoplanet detectability or specific instrument or mission capabilities; instead, it will focus on evaluating the diagnostic power of different measurements on key exoplanet science questions, simply adopting resolution, signal-to-noise, cadence, wavelength coverage as parameters along which the diagnostic power of the data will be studied. Decoupling instrumental capabilities from science goals allows this community-based effort to explore the science goals for exoplanet characterization in an unbiased manner and in a depth beyond what is possible in a typical STD T.

We envision the SAG report to be important for multiple exoplanet sub-communities and specifically foresee the following uses: 1) Future STD teams will be able to easily connect observational requirements to missions to fundamental science goals; 2) By providing an overview of the key science questions on exoplanets and how they could be answered, it may motivate new, dedicated mission proposals; 3) By providing a single, unified source of requirements on exoplanet data in advance of the Decadal Survey, the science yield of various missions designs can be evaluated realistically, with the same set of assumptions.

Our goal is to carry out this SAG study by building on both the EXOPAG and NExSS communities.

We aim to complete a report by Spring 2017 and submit it to a refereed journal, although this timeline can be adjusted to maximize the impact of the SAG15 study for the ongoing and near- future STD Ts and other mission planning processes.

Synergy with a potential future SAG proposed by Shawn Domagal-Goldman: While the SAG proposed here will include studies of habitable zone rocky planets, it will focus on planets without significant biological processes. A future SAG may be proposed by Shawn Domagal- Goldman to explore biosignatures; if such a SAG is proposed, we envision a close collaboration on these complementary, but distinct problems.

REFERENCES

- Abbot, D. S., Cowan, N. B., & Ciesla, F. J. 2012, *ApJ*, 756, 178
- Abe, Y., Abe-Ouchi, A., Sleep, N. H., & Zahnle, K. J. 2011, *Astrobiology*, 11, 443
- Ackerman, A. S. & Marley, M. S. 2001, *ApJ*, 556, 872
- Allers, K. N., Gallimore, J. F., Liu, M. C., & Dupuy, T. J. 2016, *ApJ*, 819, 133
- Allers, K. N. & Liu, M. C. 2013, *ApJ*, 772, 79
- Apai, D., Radigan, J., Buenzli, E., et al. 2013, *ApJ*, 768, 121
- Apai, D., Schneider, G., Grady, C. A., et al. 2015, *ApJ*, 800, 136
- Artigau, É., Bouchard, S., Doyon, R., & Lafrenière, D. 2009, *ApJ*, 701, 1534
- Bailey, J. 2007, *Astrobiology*, 7, 320
- Ballering, N. P., Su, K. Y. L., Rieke, G. H., & Gáspár, A. 2016, *ApJ*, 823, 108
- Barman, T. S., Macintosh, B., Konopacky, Q. M., & Marois, C. 2011a, *ApJ*, 733, 65
- Barman, T. S., Macintosh, B., Konopacky, Q. M., & Marois, C. 2011b, *ApJ*, 735, L39
- Behrens, H. & Zhang, Y. 2001, *Earth and Planetary Science Letters*, 192, 363
- Biller, B. A., Vos, J., Bonavita, M., et al. 2015, *ApJ*, 813, L23
- Black, B. A., Elkins-Tanton, L. T., Rowe, M. C., & Peate, I. U. 2012, *Earth and Planetary Science Letters*, 317, 363
- Boccaletti, A., Thalmann, C., Lagrange, A.-M., et al. 2015, *Nature*, 526, 230
- Boyer, C. & Camichel, H. 1961, *Annales d’Astrophysique*, 24, 531
- Brogi, M., de Kok, R. J., Birkby, J. L., Schwarz, H., & Snellen, I. A. G. 2014, *A&A*, 565, A124
- Buenzli, E., Apai, D., Morley, C. V., et al. 2012, *ApJ*, 760, L31
- Buenzli, E., Apai, D., Radigan, J., Reid, I. N., & Fplateau, D. 2014, *ApJ*, 782, 77
- Buenzli, E., Saumon, D., Marley, M. S., et al. 2015, *ApJ*, 798, 127

- Burgasser, A. J., Geballe, T. R., Leggett, S. K., Kirkpatrick, J. D., & Golimowski, D. A. 2006, *ApJ*, 637, 1067
- Burgasser, A. J., Looper, D. L., Kirkpatrick, J. D., Cruz, K. L., & Swift, B. J. 2008, *ApJ*, 674, 451
- Burgasser, A. J., Marley, M. S., Ackerman, A. S., et al. 2002, *ApJ*, 571, L151
- Burrows, A., Sudarsky, D., & Hubeny, I. 2006, *ApJ*, 640, 1063
- Carr, J. S. & Najita, J. R. 2008, *Science*, 319, 1504
- Chauvin, G., Lagrange, A.-M., Dumas, C., et al. 2005, *A&A*, 438, L25
- Christensen, U. R. 2010, *Space Sci. Rev.*, 152, 565
- Christensen, U. R., Holzwarth, V., & Reiners, A. 2009, *Nature*, 457, 167
- Cowan, N. B. & Abbot, D. S. 2014, *ApJ*, 781, 27
- Cowan, N. B., Abbot, D. S., & Voigt, A. 2012a, *ApJ*, 752, L3
- Cowan, N. B. & Agol, E. 2011, *ApJ*, 729, 54
- Cowan, N. B., Agol, E., Meadows, V. S., et al. 2009, *ApJ*, 700, 915
- Cowan, N. B., Fuentes, P. A., & Haggard, H. M. 2013, *MNRAS*, 434, 2465
- Cowan, N. B., Machalek, P., Croll, B., et al. 2012b, *ApJ*, 747, 82
- Cowan, N. B., Robinson, T., Livengood, T. A., et al. 2011, *ApJ*, 731, 76
- Cowan, N. B. & Strait, T. E. 2013, *ApJ*, 765, L17
- Crossfield, I. J. M., Hansen, B. M. S., Harrington, J., et al. 2010, *ApJ*, 723, 1436
- Cushing, M. C., Roellig, T. L., Marley, M. S., et al. 2006, *ApJ*, 648, 614
- de Kok, R. J., Stam, D. M., & Karalidi, T. 2011, *ApJ*, 741, 59
- de Wit, J., Gillon, M., Demory, B.-O., & Seager, S. 2012, *A&A*, 548, A128
- Debes, J. H., Weinberger, A. J., & Schneider, G. 2008, *ApJ*, 673, L191
- Demory, B.-O., de Wit, J., Lewis, N., et al. 2013, *ApJ*, 776, L25
- Dent, W. R. F., Wyatt, M. C., Roberge, A., et al. 2014, *Science*, 343, 1490

- Dupuy, T. J. & Liu, M. C. 2012, *ApJS*, 201, 19
- Faherty, J. K., Rice, E. L., Cruz, K. L., Mamajek, E. E., & Núñez, A. 2013, *AJ*, 145, 2
- Faherty, J. K., Riedel, A. R., Cruz, K. L., et al. 2016, *ApJS*, 225, 10
- Faherty, J. K., Tinney, C. G., Skemer, A., & Monson, A. J. 2014, *ApJ*, 793, L16
- Farihi, J. 2016, *New A Rev.*, 71, 9
- Foley, B. J. & Driscoll, P. E. 2016, *Geochemistry, Geophysics, Geosystems*, 17, 1885
- Ford, E. B., Seager, S., & Turner, E. L. 2001, *Nature*, 412, 885
- Fortney, J. J., Lodders, K., Marley, M. S., & Freedman, R. S. 2008, *ApJ*, 678, 1419
- Fraine, J., Deming, D., Benneke, B., et al. 2014, *Nature*, 513, 526
- Fujii, Y. & Kawahara, H. 2012, *ApJ*, 755, 101
- Fujii, Y., Kawahara, H., Suto, Y., et al. 2011, *ApJ*, 738, 184
- Fujii, Y., Kawahara, H., Suto, Y., et al. 2010, *ApJ*, 715, 866
- Fujii, Y., Kimura, J., Dohm, J., & Ohtake, M. 2014, *Astrobiology*, 14, 753
- Gibson, N. P., Aigrain, S., Barstow, J. K., et al. 2013, *MNRAS*, 436, 2974
- Goloub, P., Herman, M., Chepfer, H., et al. 2000, *J. Geophys. Res.*, 105, 14
- Greco, J. P. & Burrows, A. 2015, *ApJ*, 808, 172
- Hansen, J. E. & Hovenier, J. W. 1974, *Journal of Atmospheric Sciences*, 31, 1137
- Hansen, J. E. & Travis, L. D. 1974, *Space Sci. Rev.*, 16, 527
- Haqq-Misra, J. & Kopparapu, R. K. 2015, *MNRAS*, 446, 428
- Helling, C., Ackerman, A., Allard, F., et al. 2008, *MNRAS*, 391, 1854
- Hinkel, N. R., Timmes, F. X., Young, P. A., Pagano, M. D., & Turnbull, M. C. 2014, *AJ*, 148, 54
- Hu, R., Demory, B.-O., Seager, S., Lewis, N., & Showman, A. P. 2015, *ApJ*, 802, 51
- Hu, R., Ehlmann, B. L., & Seager, S. 2012a, *ApJ*, 752, 7

- Hu, R. & Seager, S. 2014, *ApJ*, 784, 63
- Hu, R., Seager, S., & Bains, W. 2012b, *ApJ*, 761, 166
- Hu, R., Seager, S., & Bains, W. 2013, *ApJ*, 769, 6
- Iyer, A. R., Swain, M. R., Zellem, R. T., et al. 2016, *ApJ*, 823, 109
- Joshi, M. M., Haberle, R. M., & Reynolds, R. T. 1997, *Icarus*, 129, 450
- Kaltenegger, L. & Sasselov, D. 2010, *ApJ*, 708, 1162
- Kao, M. M., Hallinan, G., Pineda, J. S., et al. 2016, *ApJ*, 818, 24
- Karalidi, T., Apai, D., Schneider, G., Hanson, J. R., & Pasachoff, J. M. 2015, *ApJ*, 814, 65
- Karalidi, T., Stam, D. M., & Hovenier, J. W. 2011, *A&A*, 530, A69
- Karalidi, T., Stam, D. M., & Hovenier, J. W. 2012, *A&A*, 548, A90
- Kaspi, Y. & Showman, A. P. 2015, *ApJ*, 804, 60
- Kasting, J. F., Whitmire, D. P., & Reynolds, R. T. 1993, *Icarus*, 101, 108
- Kataria, T., Showman, A. P., Fortney, J. J., Marley, M. S., & Freedman, R. S. 2014, *ApJ*, 785, 92
- Kawahara, H. 2016, *ApJ*, 822, 112
- Kawahara, H. & Fujii, Y. 2010, *ApJ*, 720, 1333
- Kawahara, H. & Fujii, Y. 2011, *ApJ*, 739, L62
- Kirkpatrick, J. D., Gelino, C. R., Cushing, M. C., et al. 2012, *ApJ*, 753, 156
- Knutson, H. A., Charbonneau, D., Allen, L. E., et al. 2007, *Nature*, 447, 183
- Knutson, H. A., Charbonneau, D., Cowan, N. B., et al. 2009, *ApJ*, 690, 822
- Knutson, H. A., Lewis, N., Fortney, J. J., et al. 2012, *ApJ*, 754, 22
- Kopparapu, R. k., Wolf, E. T., Haqq-Misra, J., et al. 2016, *ApJ*, 819, 84
- Kóspál, Á., Moór, A., Juhász, A., et al. 2013, *ApJ*, 776, 77
- Kreidberg, L., Bean, J. L., Désert, J.-M., et al. 2014, *Nature*, 505, 69

- Larsen, J. F. & Gardner, J. E. 2004, *Journal of Volcanology and Geothermal Research*, 134, 109
- Leconte, J., Forget, F., Charnay, B., Wordsworth, R., & Pottier, A. 2013, *Nature*, 504, 268
- Leconte, J., Wu, H., Menou, K., & Murray, N. 2015, *Science*, 347, 632
- Leggett, S. K., Cushing, M. C., Hardegree-Ullman, K. K., et al. 2016, *ArXiv e-prints*
- Liu, M. C., Magnier, E. A., Deacon, N. R., et al. 2013, *ApJ*, 777, L20
- Lodders, K. & Fegley, B. 2002, *Icarus*, 155, 393
- Lorenz, R., Lockwood, W., Barnes, J., Noll, K., & Turtle, E. 2009, in *AAS/Division for Planetary Sciences Meeting Abstracts*, Vol. 41, *AAS/Division for Planetary Sciences Meeting Abstracts #41*, 7.06
- Macintosh, B., Graham, J. R., Barman, T., et al. 2015, *Science*, 350, 64
- Madhusudhan, N., Crouzet, N., McCullough, P. R., Deming, D., & Hedges, C. 2014, *ApJ*, 791, L9
- Majeau, C., Agol, E., & Cowan, N. B. 2012, *ApJ*, 747, L20
- Malhotra, R. 1993, *Nature*, 365, 819
- Marley, M. S. & Robinson, T. D. 2015, *ARA&A*, 53, 279
- Marois, C., Macintosh, B., Barman, T., et al. 2008, *Science*, 322, 1348
- Mason, B. G., Pyle, D. M., & Oppenheimer, C. 2004, *Bulletin of Volcanology*, 66, 735
- Meng, H. Y. A., Su, K. Y. L., Rieke, G. H., et al. 2014, *Science*, 345, 1032
- Merlis, T. M. & Schneider, T. 2010, *Journal of Advances in Modeling Earth Systems*, 2, 13
- Metchev, S. A., Heinze, A., Apai, D., et al. 2015, *ApJ*, 799, 154
- Mishchenko, M. I., Rosenbush, V. K., Kiselev, N. N., et al. 2010, *ArXiv e-prints*
- Moór, A., Juhász, A., Kóspál, Á., et al. 2013, *ApJ*, 777, L25
- Morley, C. V., Fortney, J. J., Marley, M. S., et al. 2015, *ApJ*, 815, 110
- Morley, C. V., Marley, M. S., Fortney, J. J., et al. 2014, *ApJ*, 787, 78

- Mouillet, D., Larwood, J. D., Papaloizou, J. C. B., & Lagrange, A. M. 1997, *MNRAS*, 292, 896
- Oakley, P. H. H. & Cash, W. 2009, *ApJ*, 700, 1428
- O’Neill, C. & Lenardic, A. 2007, *Geophys. Res. Lett.*, 34, L19204
- O’Neill, C., Lenardic, A., Höink, T., & Coltice, N. 2013, *Mantle Convection and Outgassing on Terrestrial Planets*, ed. S. J. Mackwell, A. A. Simon-Miller, J. W. Harder, & M. A. Bullock, 473–486
- O’Neill, C., Lenardic, A., Weller, M., et al. 2016, *Physics of the Earth and Planetary Interiors*, 255, 80
- Pallé, E., Ford, E. B., Seager, S., Montañés-Rodríguez, P., & Vazquez, M. 2008, *ApJ*, 676, 1319
- Papale, P. & Polacci, M. 1999, *Bulletin of Volcanology*, 60, 583
- Parmentier, V., Fortney, J. J., Showman, A. P., Morley, C., & Marley, M. S. 2016, *ApJ*, 828, 22
- Parmentier, V., Showman, A. P., & Lian, Y. 2013, *A&A*, 558, A91
- Parol, F., Goloub, P., Herman, M., & Buriez, J.-C. 1995, in *Proc. SPIE*, Vol. 2311, *Atmospheric Sensing and Modelling*, ed. R. P. Santer, 171–180
- Perrin, M. D., Duchene, G., Millar-Blanchaer, M., et al. 2015, *ApJ*, 799, 182
- Popp, M., Schmidt, H., & Marotzke, J. 2016, *Nature Communications*, 7, 10627
- Radigan, J., Jayawardhana, R., Lafrenière, D., et al. 2012, *ApJ*, 750, 105
- Radigan, J., Lafrenière, D., Jayawardhana, R., & Artigau, E. 2014, *ApJ*, 793, 75
- Rauscher, E. & Kempton, E. M. R. 2014, *ApJ*, 790, 79
- Rauscher, E. & Menou, K. 2013, *ApJ*, 764, 103
- Reiners, A. & Basri, G. 2008, *ApJ*, 684, 1390
- Robinson, T. D., Meadows, V. S., & Crisp, D. 2010, *ApJ*, 721, L67
- Robock, A., Adams, T., Moore, M., Oman, L., & Stenchikov, G. 2007, *Geophys. Res. Lett.*, 34, L23710

- Rodigas, T. J., Debes, J. H., Hinz, P. M., et al. 2014, *ApJ*, 783, 21
- Rodler, F., Lopez-Morales, M., & Ribas, I. 2012, *ApJ*, 753, L25
- Rodriguez, S., Le Mouélic, S., Rannou, P., et al. 2011, *Icarus*, 216, 89
- Rogers, L. A. & Seager, S. 2010, *ApJ*, 712, 974
- Rossow, W. B., del Genio, A. D., & Eichler, T. 1990, *Journal of Atmospheric Sciences*, 47, 2053
- Salyk, C., Pontoppidan, K. M., Blake, G. A., et al. 2008, *ApJ*, 676, L49
- Saumon, D. & Marley, M. S. 2008, *ApJ*, 689, 1327
- Schlichting, H. E. & Sari, R. 2007, *ApJ*, 658, 593
- Schlichting, H. E., Sari, R., & Yalinewich, A. 2015, *Icarus*, 247, 81
- Schneider, G., Grady, C. A., Hines, D. C., et al. 2014, *AJ*, 148, 59
- Scholz, A., Kostov, V., Jayawardhana, R., & Mužić, K. 2015, *ApJ*, 809, L29
- Schwartz, J. C., Sekowski, C., Haggard, H. M., Pallé, E., & Cowan, N. B. 2016, *MNRAS*, 457, 926
- Seager, S., Whitney, B. A., & Sasselov, D. D. 2000, *ApJ*, 540, 504
- Showman, A. P., Fortney, J. J., Lian, Y., et al. 2009, *ApJ*, 699, 564
- Showman, A. P. & Guillot, T. 2002, *A&A*, 385, 166
- Showman, A. P., Lewis, N. K., & Fortney, J. J. 2015, *ApJ*, 801, 95
- Shporer, A. & Hu, R. 2015, *AJ*, 150, 112
- Simon, A. A., Rowe, J. F., Gaulme, P., et al. 2016, *ApJ*, 817, 162
- Sing, D. K., Fortney, J. J., Nikolov, N., et al. 2016, *Nature*, 529, 59
- Sing, D. K., Wakeford, H. R., Showman, A. P., et al. 2015, *MNRAS*, 446, 2428
- Skemer, A. J., Hinz, P. M., Esposito, S., et al. 2012, *ApJ*, 753, 14
- Skemer, A. J., Morley, C. V., Allers, K. N., et al. 2016, *ApJ*, 826, L17
- Snellen, I. A. G., Brandl, B. R., de Kok, R. J., et al. 2014, *Nature*, 509, 63

- Stam, D. M. 2008, *A&A*, 482, 989
- Stark, C. C. & Kuchner, M. J. 2008, *ApJ*, 686, 637
- Sterzik, M. F., Bagnulo, S., & Pallé, E. 2012, *Nature*, 483, 64
- Stevenson, K. B. 2016, *ApJ*, 817, L16
- Telesco, C. M., Fisher, R. S., Wyatt, M. C., et al. 2005, *Nature*, 433, 133
- Tingley, M. P., Stine, A. R., & Huybers, P. 2014, *Geophys. Res. Lett.*, 41, 7838
- Traub, W. A. & Carleton, N. P. 1975, *Journal of Atmospheric Sciences*, 32, 1045
- Tremaine, S. 1991, *Icarus*, 89, 85
- Tsiganis, K., Gomes, R., Morbidelli, A., & Levison, H. F. 2005, *Nature*, 435, 459
- Turtle, E. P., Del Genio, A. D., Barbara, J. M., et al. 2011, *Geophys. Res. Lett.*, 38, L03203
- Valencia, D., O’Connell, R. J., & Sasselov, D. D. 2007, *ApJ*, 670, L45
- van Heck, H. J. & Tackley, P. J. 2011, *Earth and Planetary Science Letters*, 310, 252
- Wagner, K., Apai, D., Kasper, M., et al. 2016, *Science*, 353, 673
- Way, M. J., Del Genio, A. D., Kiang, N. Y., et al. 2016, *Geophys. Res. Lett.*, 43, 8376
- Williams, D. M. & Gaidos, E. 2008, *Icarus*, 195, 927
- Williams, G. P. & Holloway, J. L. 1982, *Nature*, 297, 295
- Wilson, L. 1980, *Journal of Volcanology and Geothermal Research*, 8, 297
- Wolf, E. T. & Toon, O. B. 2014, *Astrobiology*, 14, 241
- Wolf, E. T. & Toon, O. B. 2015, *Journal of Geophysical Research (Atmospheres)*, 120, 5775
- Wordsworth, R. D., Forget, F., Selsis, F., et al. 2011, *ApJ*, 733, L48
- Wyatt, M. C. 2003, *ApJ*, 598, 1321
- Wyatt, M. C. 2008, *ARA&A*, 46, 339
- Wyatt, M. C., Dermott, S. F., Telesco, C. M., et al. 1999, *ApJ*, 527, 918
- Yang, H., Apai, D., Marley, M. S., et al. 2015, *ApJ*, 798, L13

- Yang, J., Boué, G., Fabrycky, D. C., & Abbot, D. S. 2014, *ApJ*, 787, L2
- Yang, J., Cowan, N. B., & Abbot, D. S. 2013, *ApJ*, 771, L45
- Zhang, X. & Showman, A. P. 2014, *ApJ*, 788, L6
- Zhou, Y., Apai, D., Schneider, G. H., Marley, M. S., & Showman, A. P. 2016, *ApJ*, 818, 176
- Zugger, M. E., Kasting, J. F., Williams, D. M., Kane, T. J., & Philbrick, C. R. 2010, *ApJ*, 723, 1168
- Zugger, M. E., Kasting, J. F., Williams, D. M., Kane, T. J., & Philbrick, C. R. 2011, *ApJ*, 739, 12